For Research Use Only

CYP11A1 Polyclonal antibody

Catalog Number: 13363-1-AP 66 Publications



Basic Information

Catalog Number: GenBank Accession Number:

13363-1-AP BC032329 GeneID (NCBI): Size:

150ul, Concentration: 450 ug/ml by 1583 Nanodrop and 300 ug/ml by Bradford $\,$ UNIPROT ID: method using BSA as the standard; P05108

Source: Full Name:

Rabbit cytochrome P450, family 11, Isotype: subfamily A, polypeptide 1

Calculated MW: Immunogen Catalog Number: 521 aa. 60 kDa AG4182 Observed MW:

49-52 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA

Cited Applications: WB, IHC, IF

Species Specificity: human, mouse, rat

Cited Species:

human, mouse, rat, pig, chicken

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human placenta tissue. HeLa cells, mouse ovary tissue, mouse testis tissue, rat testis tissue

Purification Method:

WB 1:1000-1:4000

IHC 1:100-1:400

IF/ICC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IHC: human placenta tissue, human testis tissue

IF/ICC: HepG2 cells,

Background Information

 ${\it CYP11A1}\ is\ also\ known\ as\ cytochrome\ P450C11A1, cytochrome\ P450scc\ and\ cytochrome\ P450,\ subfamily\ XIA.\ It\ is\ also\ known\ as\ cytochrome\ P450C11A1,\ cytochrome\ P450scc\ and\ cytochrome\ P450,\ subfamily\ XIA.\ It\ is\ also\ known\ as\ cytochrome\ P450C11A1,\ cytochrome\ P450scc\ and\ cytochrome\ P450,\ subfamily\ XIA.\ It\ is\ also\ known\ as\ cytochrome\ P450C11A1,\ cytochrome\ P450scc\ and\ cytochrome\ P450,\ subfamily\ XIA.\ It\ is\ also\ known\ as\ cytochrome\ P450C11A1,\ cytochrome\ P450scc\ and\ cytochrome\ P450,\ subfamily\ XIA.\ It\ is\ also\ cytochrome\ P450C11A1,\ cytochrome\ P450scc\ and\ cytochrome\ P450c\ and\ c$ the first enzyme of all steroidogenic pathways, its elimination results in syndromes of steroid deficiency, including electrolyte imbalance, aberrant gene regulation, and suppression of male sex organ development(PMID: 12145347). Expression of CYP11A1 has been a useful tool for monitoring the differentiation state of cells from varius endocrine tissues. Identification of CYP11A1 expression by 13363-1-AP detected a ~45kd band in human placenta, with respect to the reported band at 49-52kd by Roby, et al and Durkee, et al. The murine 32-kDa isoform of Cyp11a1 is not produced by omission of the first ATG-containing exon (as is the case for human CYP11A1) but perhaps by alternative translational initiation (PMID:19342447).

Notable Publications

Author	Pubmed ID	Journal	Application
Yirong Wang	36063102	Pharm Biol	WB
Shumin Li	34487832	J Steroid Biochem Mol Biol	WB
Jinzhao Zhou	35999762	Ecotoxicol Environ Saf	WB

Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13363-1-AP (CYP11A1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 13363-1-AP (CYP11A1 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 13363-1-AP (CYP11A1 Antibody) at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CYP11A1 antibody (13363-1-AP) at dilution of 1:200 and Multi-rAb Coralite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).