For Research Use Only

SPA17 Polyclonal antibody

Catalog Number: 13367-1-AP 6 Publications





Basic Information	Catalog Number: 13367-1-AP	GenBank Accession Number: BC032457	Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):	Recommended Dilutions:
	150ul , Concentration: 750 ug/ml by	53340	WB: 1:500-1:2400
	Nanodrop and 407 ug/ml by Bradford	UNIPROT ID:	IP: 0.5-4.0 ug for 1.0-3.0 mg of total
	method using BSA as the standard;	Q15506	protein lysate IHC: 1:20-1:200
	Source: Rabbit	Full Name:	IIIC. 1.20-1.200
		sperin autoantigenic protein 17	
	IgG	Calculated MW: 151 aa, 17 kDa	
	- Immunogen Catalog Number:	Observed MW:	
	AG4186	25 kDa	
Applications	Tested Applications: Positive Controls:		
	WB, IP, IHC, ELISA	WB: mou	se testis tissue,
	Cited Applications:	IP: mous	e testis tissue,
	WB, IHC, IF		nan testis tissue, human ovary tumor tissue
	Species Specificity: human, mouse, rat		
	Cited Species:		
	human, mouse		
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w	vely, antigen	
	buffer pH 6.0		
Background Information	buffer pH 6.0 Sperm surface protein Sp17(SPA17)is membrane protein. It is a highly cons spermatids, and reported to play a ro homodimer and localizes to the head	a 151 amino acids protein (17kD erved mammalian protein expres le in the interaction of sperm with l and tail of spermatozoa. SP17 ha	the zona pellucida. SP17 exists as a s been identified as a cancer/testis antiger
	buffer pH 6.0 Sperm surface protein Sp17(SPA17)is membrane protein. It is a highly cons spermatids, and reported to play a ro homodimer and localizes to the head and is expressed in ovarian cancer ar tumor immunotherapy.	a 151 amino acids protein (17kDi erved mammalian protein expres le in the interaction of sperm with I and tail of spermatozoa. SP17 ha I multiple myeloma. This sugges	ssed in developing spermatozoa and matur the zona pellucida. SP17 exists as a s been identified as a cancer/testis antiger sts that SP17 could be suitable as a target in
	buffer pH 6.0 Sperm surface protein Sp17(SPA17)is membrane protein. It is a highly cons spermatids, and reported to play a ro homodimer and localizes to the head and is expressed in ovarian cancer ar tumor immunotherapy. Author Put	a 151 amino acids protein (17kD) erved mammalian protein expres le in the interaction of sperm with and tail of spermatozoa. SP17 ha nd multiple myeloma. This sugges omed ID Journal	seed in developing spermatozoa and matur the zona pellucida. SP17 exists as a s been identified as a cancer/testis antiger ts that SP17 could be suitable as a target in Application
Background Information	buffer pH 6.0Sperm surface protein Sp17(SPA17)is membrane protein. It is a highly cons spermatids, and reported to play a ro homodimer and localizes to the head and is expressed in ovarian cancer ar tumor immunotherapy.AuthorPut Takujiro Homma330	a 151 amino acids protein (17kDi served mammalian protein expres le in the interaction of sperm with l and tail of spermatozoa. SP17 ha nd multiple myeloma. This sugges omed ID Journal 087733 Sci Rep	seed in developing spermatozoa and matur the zona pellucida. SP17 exists as a s been identified as a cancer/testis antiger sts that SP17 could be suitable as a target in Application WB
	buffer pH 6.0Sperm surface protein Sp17(SPA17)is membrane protein. It is a highly cons spermatids, and reported to play a ro homodimer and localizes to the head and is expressed in ovarian cancer ar tumor immunotherapy.AuthorPut Takujiro HommaTakujiro Homma330 30Laurie L Brunette303	a 151 amino acids protein (17kDa erved mammalian protein expres le in the interaction of sperm with and tail of spermatozoa. SP17 ha ad multiple myeloma. This sugges omed ID Journal 087733 Sci Rep 509325 BMC Cancer	seed in developing spermatozoa and matur the zona pellucida. SP17 exists as a s been identified as a cancer/testis antiger sts that SP17 could be suitable as a target in Application WB IHC
	buffer pH 6.0Sperm surface protein Sp17(SPA17)is membrane protein. It is a highly cons spermatids, and reported to play a ro homodimer and localizes to the head and is expressed in ovarian cancer ar tumor immunotherapy.AuthorPut Takujiro HommaTakujiro Homma330 30Laurie L Brunette303	a 151 amino acids protein (17kDi served mammalian protein expres le in the interaction of sperm with l and tail of spermatozoa. SP17 ha nd multiple myeloma. This sugges omed ID Journal 087733 Sci Rep	seed in developing spermatozoa and matur the zona pellucida. SP17 exists as a s been identified as a cancer/testis antiger sts that SP17 could be suitable as a target i Application WB
	buffer pH 6.0Sperm surface protein Sp17(SPA17)is membrane protein. It is a highly cons spermatids, and reported to play a ro homodimer and localizes to the head and is expressed in ovarian cancer ar tumor immunotherapy.AuthorPut Takujiro HommaTakujiro Homma330 30Laurie L Brunette303	a 151 amino acids protein (17kD) erved mammalian protein expres le in the interaction of sperm with l and tail of spermatozoa. SP17 ha ad multiple myeloma. This sugges omed ID Journal 087733 Sci Rep 309325 BMC Cancer 392314 Front Immunol er shipment.	seed in developing spermatozoa and matur the zona pellucida. SP17 exists as a s been identified as a cancer/testis antiger sts that SP17 could be suitable as a target i Application WB IHC
Notable Publications	buffer pH 6.0 Sperm surface protein Sp17(SPA17)is membrane protein. It is a highly cons spermatids, and reported to play a ro homodimer and localizes to the head and is expressed in ovarian cancer ar tumor immunotherapy. Author Put Takujiro Homma 330 Laurie L Brunette 303 Zewei Tu 355 Storage: Storage Buffer:	a 151 amino acids protein (17kDa erved mammalian protein expres le in the interaction of sperm with l and tail of spermatozoa. SP17 ha ad multiple myeloma. This sugges omed ID Journal 087733 Sci Rep 309325 BMC Cancer 392314 Front Immunol er shipment.	seed in developing spermatozoa and matur the zona pellucida. SP17 exists as a s been identified as a cancer/testis antiger sts that SP17 could be suitable as a target i Application WB IHC

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data






mouse testis tissue were subjected to SDS PAGE followed by western blot with 13367-1-AP (SPA17 antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded human testis using 13367-1-AP (SPA17 antibody) at dilution of 1:50 (under 40x lens).

Immunohistochemical analysis of paraffinembedded human testis using 13367-1-AP (SPA17 antibody) at dilution of 1:50 (under 10x lens).



IP result of anti-SPA17 (IP:13367-1-AP, 3ug; Detection:13367-1-AP 1:400) with mouse testis tissue lysate 4800ug.