For Research Use Only

Collagen Type I Polyclonal antibody



Catalog Number: 14695-1-AP

346 Publications

Basic Information

Catalog Number: 14695-1-AP

BC054498 GeneID (NCBI): Antigen affinity purification

Size:

Recommended Dilutions: WB 1:1000-1:4000

IHC 1:500-1:2000

Purification Method:

150ul , Concentration: 500 µg/ml by Nanodrop and 320 μ g/ml by Bradford Full Name: method using BSA as the standard;

collagen, type I, alpha 2

Source: Calculated MW: Rabbit 1366 aa, 130 kDa Isotype Observed MW:

Immunogen Catalog Number:

GenBank Accession Number:

AG6281

120-130 kDa

Applications

Tested Applications:

IHC. WB.FIISA

Cited Applications: IF, IHC, IP, WB

Species Specificity:

mouse, human, pig, rat

bovine, canine, human, mouse, pig, rabbit, rat, toad

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

WB: pig skin tissue, mouse lung tissue, rat lung tissue, human placenta tissue, human kidney tissue

IHC: human skin cancer tissue, human colon tissue, human oesophagus cancer tissue

Background Information

Type I collagen, the major structural component of connective tissues such as skin, tendon and bone, is the most abundant and widely expressed collagen in humans (PMID: 7620364; 8645190; 9016532). Type I collagen is a heterotrimer comprising one alpha 2(I) and two alpha 1(I) chains which are encoded by the unlinked loci COL1A2 and COL1A1 respectively. Type I collagen has a molecular mass of about 250-300 kDa, while the alpha 2(I) chain has a molecular weight of about 100-140 kDa. Mutations in COL1A2 gene are associated with osteogenesis imperfecta, Ehlers-Danlos syndrome, idiopathic osteoporosis, and atypical Marfan syndrome. This antibody raised against 1017-1366 aa of human pro-alpha 2 chain of type I collagen can recognize collagen alpha 2(I) chain and Cterminal propeptide of pro-alpha 2(I) chain.

Notable Publications

Author	Pubmed ID	Journal	Application
Lishu Feng	27689865	Angle Orthod	WB
Wenchao Gu	31557398	J Cell Mol Med	WB,IF
Yu Sun	34562065	J Cell Mol Med	WB

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



pig skin tissue were subjected to SDS PAGE followed by western blot with 14695-1-AP (Collagen Type I antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human skin cancer tissue slide using 14695-1-AP (Collagen Type I antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human skin cancer tissue slide using 14695-1-AP (Collagen Type I antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).