For Research Use Only

NUP85 Polyclonal antibody

Catalog Number: 15027-1-AP



Basic Information

Catalog Number:

15027-1-AP

GenBank Accession Number:

BC000697

GeneID (NCBI):

Size: 150ul , Concentration: 600 ug/ml by 79902

Nanodrop: **UNIPROT ID:**

Q9BW27

Rabbit Full Name:

Isotype: nucleoporin 85kDa IgG Calculated MW:

Immunogen Catalog Number: 75 kDa

AG7158 Observed MW:

60 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000 IHC 1:250-1:1000 IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: COLO 320 cells, HepG2 cells

IHC: rat brain tissue, human stomach cancer tissue,

mouse brain tissue, mouse testis tissue

IF/ICC: U2OS cells, MCF-7 cells

Background Information

This gene encodes a protein component of the Nup107-160 subunit of the nuclear pore complex. NPCs are key for nucleocytoplasmic transport, but also regulate the mitotic machinery, transcription and chromatin organization through transport-independent mechanisms. The NUP107-160 complex is acknowledged to contribute to the assembly and maintenance of the NPC structure. Members of this largest subcomplex of the NPC associate with $kine to chores, mitotic spindles, centrosomes \ and \ mitotic \ checkpoint \ regulators \ for \ proper \ completion \ of \ the \ cell$ cycle. Downregulation of NUP107-160 subcomplex members resulted in defective cytokinesis, compromised microtubule structures, altered cytoskeletal dynamics, impaired chromosome segregation and differentiation. (PMID:34170319)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 15027-1-AP (NUP85 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 15027-1-AP (NUP85 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 15027-1-AP (NUP85 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed U2OS cells using NUP85 antibody (15027-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).