For Research Use Only

HYAL2 Polyclonal antibody Catalog Number:15115-1-AP Featured Product

Featured Product 6 Publications



Basic Information	Catalog Number: 15115-1-AP	GenBank Accession BC000692	Number:	Purification Method: Antigen affinity purification	
	Size:	GeneID (NCBI):		Recommended Dilutions:	
	150ul , Concentration: 627 ug/ml by			WB 1:500-1:2000	
	Bradford method using BSA as the	UNIPROT ID:		IHC 1:50-1:500	
	standard;	Q12891		IF/ICC 1:10-1:100	
	Source:	Full Name:			
	Rabbit	hyaluronoglucosam	inidase 2		
	lsotype: IgG	Calculated MW: 54 kDa			
	Immunogen Catalog Number: AG7157	Observed MW: 54-60 kDa			
Applications	Tested Applications:		Positive Cont	rols:	
	WB, IHC, IF/ICC, ELISA Cited Applications:	WB : mouse hear liver tissue		eart tissue, HeLa cells, Raji cells, mous	
	WB, IHC, IF		IHC : human lung cancer tissue,		
	Species Specificity: human, mouse, rat		IF/ICC : Hela	cells,	
	Cited Species: human, mouse				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
	Hyaluronic acid (HA) is a glycosaminoglycan that is believed to have numerous important biologic functions, including modulation of cell proliferation, migration, and differentiation, as well as the regulation of extracellular water and protein homeostasis. It is also an integral structural component of cartilage and other tissues and acts as lubricant in joints. Hyaluronidases are a family of enzymes that catalyse the degradation of HA. In humans, there are five functional hyaluronidases: HYAL1, HYAL2, HYAL3, HYAL4 and HYAL5 (also known as SPAM1 or PH-20); plus a pseudogene, HYAL6 (also known as HYALP1). HYAL2 is present in many tissues except the adult brain (PMID: 11731268). HYAL2 can be exposed to cell surfaces that bind to the plasma membrane via GPI anchors (PMID: 11731268).				
Background Information	including modulation of cell prolifer water and protein homeostasis. It is a lubricant in joints. Hyaluronidases an are five functional hyaluronidases: H pseudogene, HYAL6 (also known as H 11731268). HYAL2 can be exposed to	also an integral struct re a family of enzyme YAL1, HYAL2, HYAL3, IYALP1). HYAL2 is pres	ural component o s that catalyse th HYAL4 and HYAL sent in many tiss	s well as the regulation of extracellula of cartilage and other tissues and acts a ne degradation of HA. In humans, there 5 (also known as SPAM1 or PH-20); plus ues except the adult brain (PMID:	
	including modulation of cell prolifer water and protein homeostasis. It is a lubricant in joints. Hyaluronidases an are five functional hyaluronidases: H pseudogene, HYAL6 (also known as H 11731268). HYAL2 can be exposed to 11731268).	also an integral struct re a family of enzyme YAL1, HYAL2, HYAL3, IYALP1). HYAL2 is pre- cell surfaces that bin	ural component (s that catalyse th HYAL4 and HYAL sent in many tisss d to the plasma r	s well as the regulation of extracellula of cartilage and other tissues and acts a ue degradation of HA. In humans, there 5 (also known as SPAM1 or PH-20); plus ues except the adult brain (PMID: nembrane via GPI anchors (PMID:	
	including modulation of cell prolifer water and protein homeostasis. It is a lubricant in joints. Hyaluronidases and are five functional hyaluronidases: H pseudogene, HYAL6 (also known as H 11731268). HYAL2 can be exposed to 11731268). Author Pu	also an integral struct e a family of enzyme YAL1, HYAL2, HYAL3, IYALP1). HYAL2 is pre- cell surfaces that bin	ural component of s that catalyse th HYAL4 and HYAL sent in many tiss d to the plasma r	s well as the regulation of extracellula of cartilage and other tissues and acts a le degradation of HA. In humans, there 5 (also known as SPAM1 or PH-20); plus ues except the adult brain (PMID: nembrane via GPI anchors (PMID:	
	including modulation of cell prolifer water and protein homeostasis. It is a lubricant in joints. Hyaluronidases and are five functional hyaluronidases: H pseudogene, HYAL6 (also known as H 11731268). HYAL2 can be exposed to 11731268). Author Pt James Fasham 34	also an integral struct e a family of enzyme YAL1, HYAL2, HYAL3, IYALP1). HYAL2 is pre- cell surfaces that bin Ibmed ID Jou 1906488 Ge	ural component of s that catalyse th HYAL4 and HYAL sent in many tisss d to the plasma r Irnal	s well as the regulation of extracellula of cartilage and other tissues and acts a ne degradation of HA. In humans, there 5 (also known as SPAM1 or PH-20); plus ues except the adult brain (PMID: nembrane via GPI anchors (PMID: Application WB,IF	
	including modulation of cell prolifer water and protein homeostasis. It is a lubricant in joints. Hyaluronidases and are five functional hyaluronidases: H pseudogene, HYAL6 (also known as H 11731268). HYAL2 can be exposed to 11731268). Author Pu James Fasham 34 Martina M A Muggenthaler 28	also an integral struct re a family of enzyme YAL1, HYAL2, HYAL3, IYALP1). HYAL2 is pre- cell surfaces that bin bomed ID Jou 1906488 Ge 3081210 PLc	ural component of s that catalyse th HYAL4 and HYAL sent in many tiss d to the plasma r	s well as the regulation of extracellula of cartilage and other tissues and acts a le degradation of HA. In humans, there 5 (also known as SPAM1 or PH-20); plus ues except the adult brain (PMID: nembrane via GPI anchors (PMID:	
	including modulation of cell prolifer water and protein homeostasis. It is a lubricant in joints. Hyaluronidases and are five functional hyaluronidases: H pseudogene, HYAL6 (also known as H 11731268). HYAL2 can be exposed to 11731268). Author Pu James Fasham 34 Martina M A Muggenthaler 28	also an integral struct e a family of enzyme YAL1, HYAL2, HYAL3, IYALP1). HYAL2 is pre- cell surfaces that bin bomed ID Jou 906488 Ge 3081210 PLC	ural component of s that catalyse th HYAL4 and HYAL sent in many tiss d to the plasma r urnal net Med	s well as the regulation of extracellula of cartilage and other tissues and acts a te degradation of HA. In humans, there 5 (also known as SPAM1 or PH-20); plus ues except the adult brain (PMID: nembrane via GPI anchors (PMID: Application WB,IF WB	
Notable Publications	including modulation of cell prolifer water and protein homeostasis. It is a lubricant in joints. Hyaluronidases at are five functional hyaluronidases: H pseudogene, HYAL6 (also known as H 11731268). HYAL2 can be exposed to 11731268). Author Pu James Fasham 34 Martina M A Muggenthaler 26 Keiichi Hiramoto 35 Storage: Storage Storage Buffer:	also an integral struct re a family of enzyme YAL1, HYAL2, HYAL3, IYALP1). HYAL2 is pre- cell surfaces that bin ibmed ID Jou i906488 Ge 3081210 PLC 1860472 De	ural component of s that catalyse th HYAL4 and HYAL sent in many tiss d to the plasma r urnal net Med	s well as the regulation of extracellula of cartilage and other tissues and acts a te degradation of HA. In humans, there 5 (also known as SPAM1 or PH-20); plus ues except the adult brain (PMID: nembrane via GPI anchors (PMID: Application WB,IF WB	
Notable Publications	including modulation of cell prolifer water and protein homeostasis. It is a lubricant in joints. Hyaluronidases an are five functional hyaluronidases: H pseudogene, HYAL6 (also known as H 11731268). HYAL2 can be exposed to 11731268). Author Pt James Fasham 34 Martina M A Muggenthaler 26 Keiichi Hiramoto 32 Storage: Stora at -20°C. Stable for one year aff Storage Buffer: PBS with 0.02% sodium azide and 50	also an integral struct re a family of enzyme YAL1, HYAL2, HYAL3, IYALP1). HYAL2 is pre- cell surfaces that bin ibmed ID Jou i906488 Ge 3081210 PLC 2860472 De rer shipment.	ural component of s that catalyse th HYAL4 and HYAL sent in many tiss d to the plasma r urnal net Med	s well as the regulation of extracellula of cartilage and other tissues and acts a te degradation of HA. In humans, there 5 (also known as SPAM1 or PH-20); plus ues except the adult brain (PMID: nembrane via GPI anchors (PMID: Application WB,IF WB	
Background Information Notable Publications Storage	including modulation of cell prolifer water and protein homeostasis. It is a lubricant in joints. Hyaluronidases at are five functional hyaluronidases: H pseudogene, HYAL6 (also known as H 11731268). HYAL2 can be exposed to 11731268). Author Pu James Fasham 34 Martina M A Muggenthaler 26 Keiichi Hiramoto 35 Storage: Storage Storage Buffer:	also an integral struct re a family of enzyme YAL1, HYAL2, HYAL3, IYALP1). HYAL2 is pre- cell surfaces that bin ibmed ID Jou i906488 Ge 3081210 PLC 2860472 De rer shipment.	ural component of s that catalyse th HYAL4 and HYAL sent in many tiss d to the plasma r urnal net Med	s well as the regulation of extracellu of cartilage and other tissues and acts the degradation of HA. In humans, ther 5 (also known as SPAM1 or PH-20); plu ues except the adult brain (PMID: nembrane via GPI anchors (PMID: Application WB,IF WB	

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 15115-1-AP (HYAL2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of Hela cells, using HYAL2 antibody 15115-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 15115-1-AP (HYAL2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 15115-1-AP (HYAL2 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).