## For Research Use Only

## KAP1 Polyclonal antibody

Catalog Number: 15202-1-AP

**Featured Product** 

29 Publications

BC004978



**Purification Method:** 

WB 1:1000-1:4000

IHC 1:1000-1:4000

protein lysate

IF 1:10-1:100

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

**Basic Information** 

**Applications** 

Catalog Number: GenBank Accession Number:

ze: GeneID (NCBI):

150ul , Concentration: 500 µg/ml by 10155 Nanodrop; Full Name

Source: tripartite motif-containing 28
Rabbit Calculated MW:

Isotype: 89 kDa
IgG Observed MW:
Immunogen Catalog Number: 100 kDa

AG7350

15202-1-AP

Tested Applications: Positive Controls:

FC, IF, IHC, IP, WB, ELISA WB: HeLa cells, rat spleen tissue, mouse testis tissue

Cited Applications: IP : HeLa cells, chIP, IF, IHC, IP, WB

Species Specificity: IHC : human lung cancer tissue, human colon tissue

human, mouse, rat IF : Hela cells,

Cited Species: human, mouse, monkey, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Background Information** 

KAP1, also named as TRIM28 or RNF96, is a 835 amino acid protein, which contain one RING-type zinc finger, one PHD-type zinc finger, one bromo domain and two B box-type zinc fingers. KAP1 localizes in the nucleus and Belongs to the TRIM/RBCC family. KAP1 is a nuclear corepressor for KRAB domain-containing zinc finger proteins and mediates gene silencing by recruiting CHD3, a subunit of the nucleosome remodeling and deacetylation (NuRD) complex, and SETDB1 to the promoter regions of KRAB target genes. KAP1 is expressed in all tissues tested including spleen, thymus, prostate, testis, ovary, small intestine, colon and peripheral blood leukocytes. The calcualted molecular weight of KAP1 is 89 kDa, but modified KAP1 is about 100 kDa. (PMID: 18590578)

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Yaguang Zhang	36100837	Sci China Life Sci	WB
Min Li	32900933	Proc Natl Acad Sci U S A	chIP
Yanhui Zhai	33539314	Reproduction	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




WB result of KAP1 antibody (15202-1-AP; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-KAP1 transfected HeLa cells.




HeLa cells were subjected to SDS PAGE followed by western blot with 15202-1-AP (KAP1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of Hela cells, using TRIM28 antibody 15202-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).




IP Result of anti-KAP1 (IP:15202-1-AP, 4ug; Detection:15202-1-AP 1:1000) with HeLa cells lysate 1200ug.




Various lysates were subjected to SDS PAGE followed by western blot with 15202-1-AP (KAP1 antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 15202-1-AP (KAP1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 15202-1-AP (KAP1 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 15202-1-AP (KAP1 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using KAP1 antibody (15202-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human KAP1 (15202-1-AP) and Coralite® 488-Conjugated Affini Pure Goat Anti-Rabbit IgG (H+L) at dilution 1:1000 (red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP, Clone:) (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set.