

For Research Use Only

CHOP; GADD153 Polyclonal antibody



Catalog Number: 15204-1-AP

Featured Product

256 Publications

Basic Information

Catalog Number:

15204-1-AP

Size:

150ul , Concentration: 700 µg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG7354

GenBank Accession Number:

BC003637

GeneID (NCBI):

1649

Full Name:

DNA-damage-inducible transcript 3

Calculated MW:

19 kDa

Observed MW:

29 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:200-1:1000

IHC 1:50-1:500

Applications

Tested Applications:

FC, IHC, WB, ELISA

Cited Applications:

CoIP, IF, IHC, WB

Species Specificity:

human, mouse

Cited Species:

Bovine, chicken, hamster, human, mouse, pig, rat, zebrafish

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : Tunicamycin treated HeLa cells, HeLa cells, MCF-7 cells

IHC : human breast cancer tissue, human cervical cancer tissue

Background Information

CHOP, also known as GADD153 or DDIT3, is a highly conserved gene in both the structural and regulatory regions. Imposed by unfolded and misfolded proteins, CHOP is significantly induced by ER stress. CHOP is considered a proapoptotic marker of ER stress dependent cell death. CHOP acts as a dominant-negative inhibitor of the transcription factor C/EBP and LAP. It may play an important role in the malignant transformation of nevus to melanoma. The calculated molecular weight of CHOP is 19 kDa, but the protein migrates on an SDS-PAGE gel with an observed molecular mass of 29 kDa (PMID: 1547942).

Notable Publications

Author	Pubmed ID	Journal	Application
Junxia Hu	31580970	Biomed Pharmacother	WB,IF
Larissa G de Vicente	34592238	Life Sci	WB
Han Liao	26415619	Chem Biol Interact	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of CHOP antibody (15204-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CHOP transfected HeLa cells.



Tunicamycin treated HeLa cells were subjected to SDS PAGE followed by western blot with 15204-1-AP (CHOP antibody at dilution of 1:300 incubated at room temperature for 1.5 hours).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 15204-1-AP (CHOP; GADD153 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10⁶ HeLa cells were stained with 0.2ug CHOP antibody (15204-1-AP, red) and control antibody (blue). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000. Cells were fixed with 4% PFA and permeabilized with 0.1% Triton X-100.