

For Research Use Only

NFKB1,p105-specific Polyclonal antibody



Catalog Number:15507-1-AP

Basic Information

Catalog Number: 15507-1-AP	GenBank Accession Number: NM_003998	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 133 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 4790	
Source: Rabbit	UNIPROT ID: P19838	
Isotype: IgG	Full Name: nuclear factor of kappa light polypeptide gene enhancer in B-cells 1	
	Calculated MW: 105 kDa	

Applications

Tested Applications:
ELISA

Species Specificity:
human, mouse, rat

Background Information

NFκB is a pleiotropic transcription factor which is present in almost all cell types and is involved in many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NFκB is activated by various intra and extra cellular stimuli such as cytokines, oxidant free radicals, ultraviolet irradiation, and bacterial or viral products. NFκB is a family of transcription factors that consists of homo and heterodimers of NFκB1/p50 and RelA/p65 subunits, and controls a variety of cellular events including development and immune responses. All members share a conserved amino terminus domain that includes dimerization, nuclear localization, and DNA binding regions, and a carboxy terminal transactivation domain. Serines 529 and 536 in the transactivation domain of RelA/p65 are phosphorylated in response to several stimuli including phorbol ester, IL1 alpha and TNF alpha as mediated by IκB kinase and p38 MAPK. Phosphorylation of serines 529 and 536 is critical for RelA/p65 transcriptional activity. Activated NFκB translocates into the nucleus and stimulates the expression of genes involved in a wide variety of biological functions. Inappropriate activation of NFκB has been associated with a number of inflammatory diseases while persistent inhibition of NFκB leads to inappropriate immune cell development or delayed cell growth. NFκB1 appears to have dual functions such as cytoplasmic retention of attached NF-κappa-B proteins by p105 and generation of p50 by a cotranslational processing. This antibody can bind p105 isoform of NFκB1 specifically.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data