

For Research Use Only

# BAG1L-specific Polyclonal ANTIBODY



Catalog Number: 16148-1-AP

1 Publications

## Basic Information

Catalog Number:

16148-1-AP

Size:

150UL, Concentration: 260 µg/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM\_004323

GeneID (NCBI):

573

Full Name:

BCL2-associated athanogene

Calculated MW:

39 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

IF 1:10-1:100

## Applications

Tested Applications:

IF, ELISA

Species Specificity:

human

Positive Controls:

IF : HepG2 cells,

## Background Information

BAG1 have been identified that modulate gene transcription through poorly defined mechanisms. Four isoforms of the BAG1 protein (BAG1S, BAG1, BAG1M and BAG1L) can be produced from a common mRNA by use of alternative translation initiation sites, including a non-canonical CTG codon in one instance. The longest, BAG1L (Mr ~50K), contains a nuclear localization signal (NLS) and resides in the nucleus, whereas BAG1M (Mr ~46K) has an incomplete NLS and distributes mainly in cytosol, unless dragged into the nucleus through interactions with other. Distribution of BAG1S(p33) is not clear yet. This antibody can recognize BAG1L and BAG1V.

## Notable Publications

Author	Pubmed ID	Journal	Application
Nada Lallous	27765852	Mol Cancer Ther	

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunofluorescent analysis of HepG2 cells, using BAG1 antibody 16148-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).