

For Research Use Only

HGD Polyclonal antibody

Catalog Number: 16465-1-AP



Basic Information

Catalog Number: 16465-1-AP	GenBank Accession Number: BC020792	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 550 µg/ml by Nanodrop;	GeneID (NCBI): 3081	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:20-1:200
Source: Rabbit	UNIPROT ID: Q93099	
Isotype: IgG	Full Name: homogentisate 1,2-dioxygenase (homogentisate oxidase)	
Immunogen Catalog Number: AG9544	Calculated MW: 37 kDa, 50 kDa	
	Observed MW: 50 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HepG2 cells, HeLa cells, rat liver tissue, mouse liver tissue

IHC : human liver tissue,

Background Information

Homogentisate 1,2-dioxygenase (HGD), also named as HGO, is a mononuclear Fe(II)-dependent oxygenase that catalyzes the third step in the pathway for the catabolism of tyrosine, the conversion of homogentisate (HG) to maleylacetoacetate (MAA) and it can exist as a dimer or trimer (PMID:14678794). HGD consists of a single type of subunit with no intermolecular disulfide bridges and requires Fe²⁺ as a cofactor (PMID: 7705358). Defects in HGD are the cause of alkaptonuria (AKU) (PMID:10594001).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 16465-1-AP (HGD antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver using 16465-1-AP (HGD antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human liver using 16465-1-AP (HGD antibody) at dilution of 1:100 (under 40x lens).