For Research Use Only

Cytokeratin 7-specific Polyclonal antibody

Catalog Number: 16857-1-AP



Basic Information

Catalog Number:

16857-1-AP NM_005556

GeneID (NCBI):

150ul , Concentration: 800 ug/ml by Nanodrop:

UNIPROT ID: Source: P08729 Rabbit Full Name: Isotype: keratin 7 IgG

> 51 kDa Observed MW:

Calculated MW:

GenBank Accession Number:

51 kDa

Applications

Tested Applications: WB, IHC, ELISA

Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:500-1:1000 IHC: 1:400-1:1600

Positive Controls:

WB: HepG2 cells,

IHC: human lung cancer tissue, human breast cancer

Background Information

Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral courterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. KRT7, also named as cytokeratin 7, is one member of type II basic cytokeratin. It is specifically expressed in the simple epithelia lining the cavities of the internal organs and in the gland ducts and blood vessels, and their neoplasms. KRT7 is marker of epithelial tissues, but not present in carcinomas of stratified squamous cell origin. This antibody is specifically against KRT7.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data







Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 16857-1-AP (Cytokeratin 7-specific antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



HepG2 cells were subjected to SDS PAGE followed by western blot with 16857-1-AP (KRT7-specific antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.