

For Research Use Only

RB1CC1 Polyclonal antibody

Catalog Number: 17250-1-AP

Featured Product

45 Publications



Basic Information

Catalog Number:

17250-1-AP

Size:

150UL, Concentration: 267 µg/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG10508

GenBank Accession Number:

BC017556

GeneID (NCBI):

9821

Full Name:

RB1-inducible coiled-coil 1

Calculated MW:

1594 aa, 183 kDa

Observed MW:

200 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:6000

IP 0.5-4.0 µg for IP and 1:500-1:1000 for WB

IHC 1:20-1:200

IF 1:10-1:100

Applications

Tested Applications:

IF, IHC, IP, WB, ELISA

Cited Applications:

co-IP, IF, IHC, IP, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, monkey, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, Jurkat cells, K-562 cells, MCF-7 cells

IP: HEK-293 cells,

IHC: human liver cancer tissue, human breast cancer tissue

IF: HepG2 cells, MCF-7 cells

Background Information

RB1CC1, also named as RBICC or FIP200, is implicated in the regulation of RB1 expression and functions as a DNA-binding transcription factor. It is a potent regulator of the RB1 pathway and a mediator that plays a crucial role in muscular differentiation. Its expression is, thus, a prerequisite for myogenic differentiation. Involved in autophagy. RB1CC1 is required for autophagosome formation. It is probably involved in the tumorigenesis of breast cancer. RB1CC1 is frequently mutated in breast cancer and shows characteristics of a classical tumor suppressor gene. This antibody is a rabbit polyclonal antibody raised against residues near the C terminus of human RB1CC1. the calculated molecular weight of RB1CC1 is 180 kDa, but the modified RB1CC1 is about 200 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Luis Muniz-Feliciano	28933590	Autophagy	WB
Le Li	30237496	Cell Death Dis	IF
Nora Wallot-Hieke	29173006	Autophagy	WB,IF,IP

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




WB result of RB1CC1 antibody (17250-1-AP, 1:2000) with si-Control and si-RB1CC1 transfected HEK293 cells.


HEK-293 cells were subjected to SDS PAGE followed by western blot with 17250-1-AP (RB1CC1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells, using RB1CC1 antibody 17250-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



IP Result of anti-RB1CC1 (IP:17250-1-AP, 5ug; Detection:17250-1-AP 1:500) with HEK-293 cells lysate 2500ug.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 17250-1-AP (RB1CC1 Antibody) at dilution of 1:50 (under 40x lens).