For Research Use Only

ACADL-Specific Polyclonal antibody

Catalog Number: 17526-1-AP

Featured Product

58 Publications



Basic Information

Catalog Number:

17526-1-AP

Size:

NM 001608

GeneID (NCBI):

GenBank Accession Number:

150ul , Concentration: 500 ug/ml by

Nanodrop: Source:

UNIPROT ID: P28330 Full Name:

Rabbit Isotype: IgG

acyl-Coenzyme A dehydrogenase,

long chain

Calculated MW: 48 kDa

Observed MW:

48 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions:

WB: 1:5000-1:50000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

Applications

Tested Applications:

WB, IP, ELISA

Cited Applications:

WB, IHC, IF, IP Species Specificity:

human, mouse

Cited Species:

human, mouse, rat, pig, sheep, ewes

Positive Controls:

WB: mouse kidney tissue, mouse heart tissue, mouse

liver tissue, mouse lung tissue

IP: mouse kidney tissue,

Background Information

ACADL(Long-chain specific acyl-CoA dehydrogenase, mitochondrial) is also named as LCAD and belongs to the acyl- $CoA\ dehydrogenase\ family. It\ catalyzes\ the\ first\ reaction\ of\ the\ mitochondrial\ \beta-oxidation\ of\ fatty\ acids\ and\ is$ synthesized in the cytosol as a precursor that is larger than its mature from. The size difference between the precursor and mature rat LCAD has been estimated to be 3 kDa on the basis of SDS-PAGE(PMID:1774065). Defects in ACADL are a cause of acyl-CoA dehydrogenase very long-chain deficiency (ACADVLD). This antibody is specific to ACADL.

Notable Publications

Author	Pubmed ID	Journal	Application
Tingting Lv	33659423	Bio Protoc	WB, IP
De Huang	25242319	Cell Rep	WB, IHC
Dharendra Thapa	30323061	J Biol Chem	WB

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 17526-1-AP (ACADL-Specific antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



IP result of anti-ACADL-Specific (IP:17526-1-AP, 4ug; Detection:17526-1-AP 1:10000) with mouse kidney tissue lysate 2160 ug.