### For Research Use Only

# HMGCS1 Polyclonal antibody

Catalog Number: 17643-1-AP

Featured Product

40 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

 17643-1-AP
 BC083514

 Size:
 GeneID (NCBI):

 150ul , Concentration: 700 ug/ml by
 3157

Nanodrop; UNIPROT ID:
Source: Q01581
Rabbit Full Name:

 Isotype:
 3-hydroxy-3-methylglutaryl 

 IgG
 Coenzyme A synthase 1 (soluble)

Immunogen Catalog Number:Calculated MW:AG11825520 aa, 57 kDa

Observed MW: 57 kDa Purification Method: Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate
IHC 1:300-1:1200
IF/ICC 1:400-1:1600

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, IP, CoIP Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:** 

WB: A431 cells, A549 cells, HepG2 cells, LNCaP cells,

mouse liver tissue

IP: mouse liver tissue,

IHC: human ovary cancer tissue,

IF/ICC: HepG2 cells,

# **Background Information**

HMGCS1(Hydroxymethylglutaryl-CoA synthase, cytoplasmic) is also named as HMGCS and belongs to the HMG-CoA synthase family. The gene encodes a 520 amino acid protein which catalyzes the condensation of acetyl-CoA and acetoacetyl-CoA into 3-hydroxy-3-methylglutaryl CoA and participates in the mevalonate pathway but its end product is cholesterol(PMID:20346956). The human and hamster enzymes are distinguished by differential inhibition by magnesium ion(PMID:2870496).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Jiang-Yun Peng	36307901	Adv Sci (Weinh)	WB
Feng Zhao	34803510	Int J Biol Sci	WB
Catherine I Real	24513310	Antiviral Res	WB

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




WB result of HMGCS1 antibody (17643-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HMGCS1 transfected HepG2 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 17643-1-AP (HMGCS1 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 17643-1-AP (HMGCS1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-HMGCS1 (IP:17643-1-AP, 3ug; Detection:17643-1-AP 1:1000) with mouse liver tissue lysate 4000ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using HMGCS1 antibody (17643-1-AP) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 17643-1-AP (HMGCS1 antibody) at dilution of 1:600 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).