

For Research Use Only

DHX9 Polyclonal antibody

Catalog Number: 17721-1-AP

Featured Product

42 Publications



Basic Information

Catalog Number:

17721-1-AP

Size:

150ul, Concentration: 700 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG12104

GenBank Accession Number:

BC014246

GeneID (NCBI):

1660

UNIPROT ID:

Q08211

Full Name:

DEAH (Asp-Glu-Ala-His) box polypeptide 9

Calculated MW:

1270 aa, 141 kDa

Observed MW:

140 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:5000-1:50000

IHC: 1:250-1:1000

IF/ICC: 1:400-1:1600

FC (Intra): 0.25 ug per 10⁶ cells in a 100 µl suspension

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP, ChIP, RIP, ELISA

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, mouse testis tissue, rat spleen tissue, rat testis tissue, Jurkat cells

IHC: mouse brain tissue, mouse testis tissue

IF/ICC: HepG2 cells,

FC (Intra): HepG2 cells,

Background Information

RNA helicases play important roles in transcription, RNA processing, translation, and RNA replication. DEAD box proteins are putative RNA helicases that have a characteristic Asp-Glu-Ala-Asp (DEAD) box as 1 of 8 highly conserved sequence motifs. DHX9 a member of the DEAH family of proteins, which possess a double-stranded RNA-binding domain (dsRBD) and a helicase domain [PMID:20569003]. It unwinds double-stranded DNA and RNA in a 3' to 5' direction. Alteration of secondary structure of DHX9 may subsequently influence interactions with proteins or other nucleic acids. It is also a component of the CRD-mediated complex that promotes MYC mRNA stability. In addition, it is involved with LARP6 in the stabilization of type I collagen mRNAs for CO1A1 and CO1A2 [PMID:19029303, 22190748].

Notable Publications

Author	Pubmed ID	Journal	Application
Feng Shi	34676915	J Clin Lab Anal	WB, IHC
Liang Liu	36377508	CNS Neurosci Ther	WB, IHC
Nila Roy Choudhury	29117863	BMC Biol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 17721-1-AP (DHX9 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 17721-1-AP (DHX9 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 17721-1-AP (DHX9 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using DHX9 antibody (17721-1-AP) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using DHX9 antibody (17721-1-AP) at dilution of 1:800 and Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-Phalloidin (red).



1x10⁶ HepG2 cells were intracellularly stained with 0.25 ug DHX9 Polyclonal antibody (17721-1-AP) and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).