

For Research Use Only

# PD-L1/CD274 Polyclonal antibody

Catalog Number: 17952-1-AP

Featured Product

199 Publications



## Basic Information

### Catalog Number:

17952-1-AP

### Size:

150ul, Concentration: 700 ug/ml by Nanodrop;

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG12432

### GenBank Accession Number:

BC074984

### GeneID (NCBI):

29126

### UNIPROT ID:

Q9NZQ7

### Full Name:

CD274 molecule

### Calculated MW:

290 aa, 33 kDa

### Observed MW:

45-56 kDa, 65-70 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB: 1:500-1:1000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC: 1:300-1:1200

IF/ICC: 1:10-1:100

## Applications

### Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

### Cited Applications:

WB, IHC, IF, IP, CoIP, CHIP

### Species Specificity:

human, mouse, rat

### Cited Species:

human, mouse, rat

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB:** human placenta tissue, A375 cells, IFN gamma treated A549 cells, HeLa cells, HepG2 cells, human heart tissue, K-562 cells, mouse heart tissue, mouse skeletal muscle tissue, rat kidney tissue

**IP:** mouse heart tissue,

**IHC:** human tonsillitis tissue, mouse heart tissue, human stomach cancer tissue

**IF/ICC:** HEK-293 cells,

## Background Information

Programmed cell death ligand 1 (PD-L1, CD274, or B7-H1), is the first member of B7 family to be discovered. B7 family molecules are type I transmembrane proteins belonging to the immunoglobulin superfamily. In concert with their CD28 family receptors, the B7s are key regulators of the adaptive immune response. PD-L1 is suggested as a negative regulator of T and B cell, and plays important role in mediating tolerance of lymphocytes to self-antigens. It is also involved in the costimulatory signal, essential for T-cell proliferation and production of IL10 and IFNG, in an IL2-dependent and a PD-1-independent manner. PD-L1 is a 290 aa transmembrane protein with a calculated molecular weight of 33 kDa, it is predicted to be 27-30 kDa after signal peptide cleavage (PMID: 25609200; 17076679). The apparent molecular weight has also been reported as 45-70 kDa, major glycosylated form of 45-50 kDa and multiple post-translational modifications form of 65-70 kDa (PMID: 18760278; 16493058).

## Notable Publications

Author	Pubmed ID	Journal	Application
Ching-Yao Yang	29036791	Cancer Biomark	IHC
Gagan Chhabra	34597611	J Invest Dermatol	IHC
Haoyu Guo	36201949	Biomaterials	IHC

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Untreated and IFN gamma treated A549 cells were subjected to SDS PAGE followed by western blot with 17952-1-AP (PD-L1/CD274 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




human placenta tissue were subjected to SDS PAGE followed by western blot with 17952-1-AP (PD-L1/CD274 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 17952-1-AP (PD-L1/CD274 antibody) at dilution of 1:600 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 17952-1-AP (PD-L1/CD274 antibody) at dilution of 1:600 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HEK-293 cells using 17952-1-AP (PD-L1/CD274 antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP result of anti-PD-L1/CD274 (IP:17952-1-AP, 4ug; Detection:66248-1-Ig 1:1000) with mouse heart tissue lysate 4000ug.