# For Research Use Only

# Cannabinoid receptor 1 Polyclonal antibody

Proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Catalog Number: 17978-1-AP

29 Publications

### **Basic Information**

Catalog Number: 17978-1-AP

GenBank Accession Number:

BC074812

Size: GenelD (NCBI):

150ul , Concentration: 500 ug/ml by 1268

Nanodrop; UNIPROT ID:
Source: P21554
Rabbit Full Name:

Isotype: cannabinoid receptor 1 (brain)

IgG Calculated MW:
Immunogen Catalog Number: 472 aa, 53 kDa
AG12371 Observed MW:

45 kDa, 66 kDa

Purification Method: Antigen affinity purification

Recommended Dilutions: WB: 1:1000-1:4000 IHC: 1:200-1:800 IF-P: 1:50-1:500

# **Applications**

Tested Applications: WB, IHC, IF-P, ELISA

Cited Applications: WB, IHC, IF

Species Specificity: human, mouse, rat Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

WB: SH-SY5Y cells, mouse brain tissue, rat brain tissue

IHC: mouse brain tissue,
IF-P: mouse brain tissue,

# **Background Information**

Cannabinoid receptor 1 (CNR1, or CB1) and CNR2 (CB2) are members of the guanine-nucleotide-binding protein (G-protein) coupled receptor family. The CB1 receptor is expressed mainly in the brain. The CB2 receptor is expressed mainly in the immune system and in hematopoietic cells. The two receptors have been found to be involved in the cannabinoid-induced CNS effects (including alterations in mood and cognition) experienced by users of marijuana.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Tingting Guo	31634502	Neurosci Lett	IHC
Kaminitz Ayelet A	24219803	World J Biol Psychiatry	
An Liu	34037653	Food Funct	WB

# Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3  $\,$ 

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


# **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 17978-1-AP (Cannabinoid receptor 1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 17978-1-AP (Cannabinoid receptor 1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 17978-1-AP (Cannabinoid receptor 1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using Cannabinoid receptor 1 antibody (17978-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 17978-1-AP (Cannabinoid receptor 1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.