

For Research Use Only

# CD41/Integrin alpha 2b Polyclonal antibody

Catalog Number: 18308-1-AP

7 Publications



## Basic Information

### Catalog Number:

18308-1-AP

### Size:

150ul, Concentration: 700 ug/ml by Nanodrop;

### Source:

Rabbit

### Isotype:

IgG

### GenBank Accession Number:

BC126442

### GeneID (NCBI):

3674

### UNIPROT ID:

P08514

### Full Name:

integrin, alpha 2b (platelet glycoprotein IIb of IIb/IIIa complex, antigen CD41)

### Calculated MW:

113 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

IHC 1:500-1:2000

## Applications

### Tested Applications:

IHC, ELISA

### Cited Applications:

WB, IF

### Species Specificity:

human, mouse, rat

### Cited Species:

human, mouse, rat

### Positive Controls:

IHC : mouse spleen tissue, human spleen tissue

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

The integrins are a superfamily of cell adhesion receptors that bind to extracellular matrix ligands, cell-surface ligands, and soluble ligands (PMID: 17543136). They are transmembrane  $\alpha\beta$  heterodimers and at least 24 distinct integrin heterodimers are formed by the combination of 18  $\alpha$  and eight  $\beta$  known subunits (PMID: 17543136; 20029421). In addition to mediating cell adhesion, integrins also play important roles in modulating signal transduction pathways that control cellular responses including migration, proliferation, differentiation, and apoptosis (PMID:19118207). Integrin alpha-IIb (ITGA2B, CD41) is expressed on platelets, megakaryocytes and some hematopoietic progenitor cells (PMID: 11934866). This protein undergoes post-translational cleavage to yield disulfide linked light (25 kDa) and heavy (125 kDa) chains which join with integrin beta3 (CD61) to form a receptor for fibronectin, fibrinogen, plasminogen, prothrombin, thrombospondin and vitronectin. Integrin alpha-IIb/beta-3 is crucial for platelet aggregation through binding of soluble fibrinogen.

## Notable Publications

Author	Pubmed ID	Journal	Application
Ling-Wei Hsu	32877952	Thromb Haemost	IF
Tianyi Wang	36195952	J Nanobiotechnology	WB
Xiaolu Zhu	30679324	Haematologica	IF

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 18308-1-AP (CD41/Integrin alpha 2b antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 18308-1-AP (CD41/Integrin alpha 2b antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).