

For Research Use Only

PSAP Polyclonal antibody

Catalog Number: 18423-1-AP

Featured Product



Basic Information

Catalog Number:

18423-1-AP

Size:

150ul, Concentration: 600 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG13266

GenBank Accession Number:

BC001503

GeneID (NCBI):

5660

UNIPROT ID:

P07602

Full Name:

prosaposin

Calculated MW:

58 kDa

Observed MW:

60-70 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IHC 1:20-1:200

Applications

Tested Applications:

WB, IHC, ELISA

Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HEK-293T cells, HeLa cells, HepG2 cells

IHC : human liver cancer tissue, human prostate cancer tissue

Background Information

proteins' (SAPs) that assist in the lysosomal hydrolysis of sphingolipids. After proteolytic processing of the prosaposin protein, these 4 released polypeptides are functional activators. Saposin A was encoded by residues 60 to 143 of PSAP, saposin B by 195 to 275, saposin C by 311 to 390, and saposin D by 405 to 487. There are four 12-14 kDa heatstable glycoproteins. This antibody should recognize saposin D.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 18423-1-AP (PSAP antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver cancer using 18423-1-AP (PSAP antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human liver cancer using 18423-1-AP (PSAP antibody) at dilution of 1:50 (under 40x lens).



WB result of PSAP antibody (18423-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PSAP transfected HepG2 cells.