For Research Use Only

nNOS Polyclonal antibody Catalog Number: 18984-1-AP 6 Publications



Basic Information	Catalog Number: 18984-1-AP	GenBank Accession Number: NM_000620	Purification Method: Antigen affinity purification
	Size: 150ul , Concentration: 800 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG	GeneID (NCBI): 4842 UNIPROT ID: P29475 Full Name: nitric oxide synthase 1 (neuro Calculated MW: 161 kDa	Recommended Dilutions: IHC 1:50-1:500 IF-P 1:50-1:500
Applications	Tested Applications:	Positive Controls: IHC : mouse heart tissue, mouse kidney tissue, rat testis tissue, mouse brain tissue	
	IHC, IF-P, ELISA Cited Applications:		
	IHC IF-P : mouse bra Species Specificity: human, mouse, rat		mouse brain tissue,
	Cited Species: human, mouse, rat, pig, canine		
	Note-IHC: suggested antigen i TE buffer pH 9.0; (*) Alternati	vely, antigen	
	retrieval may be performed w buffer pH 6.0	vith citrate	
Background Information	nNOS belongs to the NOS family. nN functions throughout the body. In the neurotransmitter. NO is a reactive fre neurotransmission and antimicrobia synthases. nNOS is a nitric oxide syn	OS produces nitric oxide (NO) brain and peripheral nervous re radical which acts as a biolo l and antitumoral activities. No thase which is highly expresse	which is a messenger molecule with diverse system, NO displays many properties of a Igic mediator in several processes, including D is synthesized from L-arginine by nitric oxide ed in skeletal muscle. Genetic variations in nNC loric stenosis type 1 (IHPS1). The antibody can
	nNOS belongs to the NOS family. nN functions throughout the body. In the neurotransmitter. NO is a reactive fre neurotransmission and antimicrobia synthases. nNOS is a nitric oxide syn gene are associated with susceptibil recognize isoform 1,2,4 of nNOS.	OS produces nitric oxide (NO) brain and peripheral nervous re radical which acts as a biolo l and antitumoral activities. No thase which is highly expresse	system, NO displays many properties of a gic mediator in several processes, including O is synthesized from L-arginine by nitric oxide ed in skeletal muscle. Genetic variations in nNO
	buffer pH 6.0 nNOS belongs to the NOS family. nN functions throughout the body. In the neurotransmitter. NO is a reactive fre neurotransmission and antimicrobia synthases. nNOS is a nitric oxide syn gene are associated with susceptibil recognize isoform 1,2,4 of nNOS.	OS produces nitric oxide (NO) brain and peripheral nervous eradical which acts as a biolo l and antitumoral activities. No thase which is highly expresse ity to infantile hypertrophic py	system, NO displays many properties of a gic mediator in several processes, including D is synthesized from L-arginine by nitric oxide ed in skeletal muscle. Genetic variations in nNO cloric stenosis type 1 (IHPS1). The antibody can Application
Background Information Notable Publications	buffer pH 6.0 nNOS belongs to the NOS family. nN functions throughout the body. In the neurotransmitter. NO is a reactive fre neurotransmission and antimicrobia synthases. nNOS is a nitric oxide syn gene are associated with susceptibil recognize isoform 1,2,4 of nNOS. Author Put Lei Wu 326	OS produces nitric oxide (NO) brain and peripheral nervous seradical which acts as a biolo l and antitumoral activities. No thase which is highly expresse ity to infantile hypertrophic py Journal	system, NO displays many properties of a gic mediator in several processes, including D is synthesized from L-arginine by nitric oxide ed in skeletal muscle. Genetic variations in nNO cloric stenosis type 1 (IHPS1). The antibody can Application
	buffer pH 6.0 nNOS belongs to the NOS family. nN functions throughout the body. In the neurotransmitter. NO is a reactive free neurotransmission and antimicrobia synthases. nNOS is a nitric oxide syn gene are associated with susceptibil recognize isoform 1,2,4 of nNOS. Author Put Lei Wu 326 Alireza Badiei 349	OS produces nitric oxide (NO) brain and peripheral nervous se radical which acts as a biolo l and antitumoral activities. No thase which is highly expresse ity to infantile hypertrophic py pomed ID Journal ACS Chem No.	system, NO displays many properties of a ogic mediator in several processes, including D is synthesized from L-arginine by nitric oxide ed in skeletal muscle. Genetic variations in nNC cloric stenosis type 1 (IHPS1). The antibody can Application eurosci
	buffer pH 6.0 nNOS belongs to the NOS family. nN functions throughout the body. In the neurotransmitter. NO is a reactive free neurotransmission and antimicrobia synthases. nNOS is a nitric oxide syn gene are associated with susceptibil recognize isoform 1,2,4 of nNOS. Author Put Lei Wu 326 Alireza Badiei 349	OS produces nitric oxide (NO) brain and peripheral nervous ser radical which acts as a biolo land antitumoral activities. No thase which is highly expresse ity to infantile hypertrophic py omed ID Journal ACS Chem No 254206 Exp Eye Res 225405 Mod Pathol er shipment.	system, NO displays many properties of a gic mediator in several processes, including D is synthesized from L-arginine by nitric oxide ed in skeletal muscle. Genetic variations in nNC cloric stenosis type 1 (IHPS1). The antibody can Application eurosci IHC

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



WWW(PfGLAB.COM

Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 18984-1-AP (nNOS antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 18984-1-AP (nNOS antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse brain tissue using nNOS antibody (18984-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).