For Research Use Only

Urocortin Polyclonal antibody

Catalog Number:19731-1-AP 1 Publications



Basic Information	Catalog Number: 19731-1-AP	GenBank Accession Number: NM_003353	Purification Method: Antigen affinity purification	
	Size: 150ul , Concentration: 500 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG	GenelD (NCBI):	Recommended Dilutions:	
		7349 UNIPROT ID: P55089	WB 1:500-1:3000	
			IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
		Full Name:		
		urocortin		
		Calculated MW: 13 kDa		
		Observed MW: 22 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IP, ELISA Cited Applications:		ouse kidney tissue, HEK-293 cells, HeLa cells, kidney tissue, human testis tissue use kidney tissue,	
	WB, IHC			
	Species Specificity: human		,	
	Cited Species:			
	human			
	Urocortin is a member of the sauvagine/corticotropin-releasing factor/urotensin I family that includes CRF, urotensin I, sauvagine, urocortin II and urocortin III. Urocortin is a potent anorexigenic peptide that induces fed-lik motor activity when administered centrally or peripherally in fasted animals. Urocortin is also a potent and long- lasting hypotensive agent and increases coronary blood flow. Urocortin acts in vitro to stimulate the secretion of adrenocorticotropic hormone (ACTH) and binds with high affinity to CRF receptor types 1, 2-alpha, and 2-beta. Urocortin is synthesized as precursor, which is subsequently processed to the mature biologically active peptide (40-amino acid Molecule). The 22 kDa may be a urocortin 1 precursor protein (PMID: 28887029).			
Background Information	motor activity when administered ce lasting hypotensive agent and increa adrenocorticotropic hormone (ACTH) Urocortin is synthesized as precursor	nd urocortin III. Urocortin is a p entrally or peripherally in faste ases coronary blood flow. Uroco and binds with high affinity to , which is subsequently proces	otent anorexigenic peptide that induces fed-lik ed animals. Urocortin is also a potent and long- ortin acts in vitro to stimulate the secretion of o CRF receptor types 1, 2-alpha, and 2-beta. sed to the mature biologically active peptide (
Background Information	motor activity when administered ce lasting hypotensive agent and increa adrenocorticotropic hormone (ACTH) Urocortin is synthesized as precursor 40-amino acid Molecule). The 22 kD	nd urocortin III. Urocortin is a p entrally or peripherally in faste ases coronary blood flow. Uroco and binds with high affinity to , which is subsequently proces	otent anorexigenic peptide that induces fed-lik ed animals. Urocortin is also a potent and long- ortin acts in vitro to stimulate the secretion of o CRF receptor types 1, 2-alpha, and 2-beta. sed to the mature biologically active peptide (a	
	advensini, sadvagine, diocordini ar motor activity when administered ce lasting hypotensive agent and increa adrenocorticotropic hormone (ACTH) Urocortin is synthesized as precursor 40-amino acid Molecule). The 22 kD	nd urocortin III. Urocortin is a p entrally or peripherally in faste ases coronary blood flow. Uroco and binds with high affinity to , which is subsequently proces a may be a urocortin 1 precurs	otent anorexigenic peptide that induces fed-lil ed animals. Urocortin is also a potent and long- ortin acts in vitro to stimulate the secretion of o CRF receptor types 1, 2-alpha, and 2-beta. sed to the mature biologically active peptide (or protein (PMID: 28887029). Application	
Notable Publications	adversifi, savvagine, diveoritini ar motor activity when administered ce lasting hypotensive agent and increa adrenocorticotropic hormone (ACTH) Urocortin is synthesized as precursor 40-amino acid Molecule). The 22 kD Author Pul Xiaolan Guo 38	nd urocortin III. Urocortin is a pentrally or peripherally in faste ases coronary blood flow. Uroco and binds with high affinity to , which is subsequently proces a may be a urocortin 1 precurs	otent anorexigenic peptide that induces fed-li ed animals. Urocortin is also a potent and long- ortin acts in vitro to stimulate the secretion of 0 CRF receptor types 1, 2-alpha, and 2-beta. sed to the mature biologically active peptide (or protein (PMID: 28887029). Application	
Notable Publications	autoerism, savvagine, diccontini ar motor activity when administered ce lasting hypotensive agent and increa adrenocorticotropic hormone (ACTH) Urocortin is synthesized as precursor 40-amino acid Molecule). The 22 kD Author Pul Xiaolan Guo 38 Storage: Storage Storage Multiple Storage Buffer:	nd urocortin III. Urocortin is a pentrally or peripherally in faste ases coronary blood flow. Uroc and binds with high affinity to , which is subsequently proces a may be a urocortin 1 precurs bmed ID Journal 546882 J Cancer Res	otent anorexigenic peptide that induces fed-li ed animals. Urocortin is also a potent and long- ortin acts in vitro to stimulate the secretion of 0 CRF receptor types 1, 2-alpha, and 2-beta. sed to the mature biologically active peptide for protein (PMID: 28887029). Application	
	adversifi, savvagine, diveoritini ar motor activity when administered ce lasting hypotensive agent and increa adrenocorticotropic hormone (ACTH) Urocortin is synthesized as precursor 40-amino acid Molecule). The 22 kD Author Pul Xiaolan Guo 38. Storage: Storage: Store at -20°C. Stable for one year affect	nd urocortin III. Urocortin is a pentrally or peripherally in faste asses coronary blood flow. Uroc and binds with high affinity to which is subsequently proces a may be a urocortin 1 precurs bmed ID Journal 546882 J Cancer Res ter shipment.	otent anorexigenic peptide that induces fed-lil ed animals. Urocortin is also a potent and long- ortin acts in vitro to stimulate the secretion of 0 CRF receptor types 1, 2-alpha, and 2-beta. sed to the mature biologically active peptide (or protein (PMID: 28887029). Application	


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



 $\begin{array}{c}
50kd \rightarrow \\
40kd \rightarrow \\
20kd \rightarrow \\
20kd \rightarrow \\
15kd \rightarrow \\
10kd \rightarrow \\
5kd \rightarrow \\
\end{array}$



Various lysates were subjected to SDS PAGE followed by western blot with 19731-1-AP (Urocortin antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours. HeLa cells were subjected to SDS PAGE followed by western blot with 19731-1-AP (Urocortin antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.

IP result of anti-Urocortin (IP:19731-1-AP, 4ug; Detection:19731-1-AP 1:1000) with mouse kidney tissue lysate 2160 ug.