For Research Use Only

Periostin Polyclonal antibody

Catalog Number:19899-1-AP

Featured Product 38 Publications



Basic Information	Catalog Number: 19899-1-AP	GenBank Accession Number: BC 106710	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 255 ug/ml by Nanodrop;	10631	IHC: 1:50-1:500	
		UNIPROT ID:		
	Source: Rabbit Isotype:	Q15063		
		Full Name: periostin, osteoblast specific factor Calculated MW:		
	IgG			
	Immunogen Catalog Number: AG13791	93 kDa		
Applications	Tested Applications: IHC, ELISA	Positive Controls:		
	Cited Applications: IHC, IF	IHC : human colon tissue, human breast cancer tiss human skin cancer tissue		
	Species Specificity: human, mouse, rat			
	Cited Species:			
	human, mouse, rat, zebrafish			
	Note-IHC: suggested antigen (TE buffer pH 9.0; (*) Alternati			
	retrieval may be performed w buffer pH 6.0			
Background Information	retrieval may be performed w buffer pH 6.0 Periostin (POSTN, PN), originally nar now classified as a matricellular pro has a role in bone, tooth and heart de broad range of human cancer types, i periostin is expressed by fibroblasts	with citrate ned as osteoblast-specific fact tein. It is present in a wide var evelopment and function. Stud ncluding lung, ovary, breast ar in the normal tissue and in the ce. The isoforms of periostin a rized by individual presence of	or 2 (OSF-2), is a 90-kDa secreted protein iety of normal adult tissues and fetal tiss ies show that periostin is overexpressed i nd colon cancers. Recent evidence reveals a stroma of the primary tumour, and it is n re between 83 and 93 kDa in mass and di or absence of cassette exons 17-21	ues, a in a s that equire
	retrieval may be performed w buffer pH 6.0 Periostin (POSTN, PN), originally nar now classified as a matricellular pro has a role in bone, tooth and heart de broad range of human cancer types, i periostin is expressed by fibroblasts to allow cancer stem cell maintenan their C-terminal sequences, character (UniProtKB/Swiss-Prot,PMID: 219977	with citrate ned as osteoblast-specific fact tein. It is present in a wide var evelopment and function. Stud ncluding lung, ovary, breast ar in the normal tissue and in the ce. The isoforms of periostin a rized by individual presence of	iety of normal adult tissues and fetal tiss ies show that periostin is overexpressed i nd colon cancers. Recent evidence reveals a stroma of the primary tumour, and it is re re between 83 and 93 kDa in mass and di	ues, ai n a s that equire ffer in
Background Information	retrieval may be performed w buffer pH 6.0 Periostin (POSTN, PN), originally nar now classified as a matricellular pro has a role in bone, tooth and heart de broad range of human cancer types, i periostin is expressed by fibroblasts to allow cancer stem cell maintenan their C-terminal sequences, character (UniProtKB/Swiss-Prot,PMID: 219977	ned as osteoblast-specific fact tein. It is present in a wide var evelopment and function. Stud ncluding lung, ovary, breast ar in the normal tissue and in the ce. The isoforms of periostin a rized by individual presence of 59).	iety of normal adult tissues and fetal tiss ies show that periostin is overexpressed i nd colon cancers. Recent evidence reveals a stroma of the primary tumour, and it is n re between 83 and 93 kDa in mass and di or absence of cassette exons 17-21 Applicatio	ues, an in a 5 that equire ffer in
	retrieval may be performed w buffer pH 6.0 Periostin (POSTN, PN), originally nar now classified as a matricellular pro has a role in bone, tooth and heart de broad range of human cancer types, i periostin is expressed by fibroblasts to allow cancer stem cell maintenan their C-terminal sequences, characte (UniProtKB/Swiss-Prot,PMID: 219977 Author Pu Han Li 32	ned as osteoblast-specific fact tein. It is present in a wide var velopment and function. Stud ncluding lung, ovary, breast ar in the normal tissue and in the ce. The isoforms of periostin a rized by individual presence of 59). bmed ID Journal 964617 J Biomed Ma	iety of normal adult tissues and fetal tiss ies show that periostin is overexpressed i nd colon cancers. Recent evidence reveals a stroma of the primary tumour, and it is n re between 83 and 93 kDa in mass and di or absence of cassette exons 17-21 Applicatio	ues, ai n a s that equire ffer in
	retrieval may be performed w buffer pH 6.0 Periostin (POSTN, PN), originally nar now classified as a matricellular pro has a role in bone, tooth and heart de broad range of human cancer types, i periostin is expressed by fibroblasts to allow cancer stem cell maintenan their C-terminal sequences, characte (UniProtKB/Swiss-Prot,PMID: 219977 Author Pu Han Li 32 Xuanwen Xu 36	ned as osteoblast-specific fact tein. It is present in a wide var velopment and function. Stud ncluding lung, ovary, breast ar in the normal tissue and in the ce. The isoforms of periostin a rized by individual presence of 59). bmed ID Journal 964617 J Biomed Ma	iety of normal adult tissues and fetal tiss ies show that periostin is overexpressed i nd colon cancers. Recent evidence reveals a stroma of the primary tumour, and it is re re between 83 and 93 kDa in mass and di or absence of cassette exons 17-21 Application ter Res A IHC	ues, a in a 5 that equire ffer in
	retrieval may be performed w buffer pH 6.0 Periostin (POSTN, PN), originally nar now classified as a matricellular pro has a role in bone, tooth and heart de broad range of human cancer types, i periostin is expressed by fibroblasts to allow cancer stem cell maintenan their C-terminal sequences, characte (UniProtKB/Swiss-Prot,PMID: 219977 Author Pu Han Li 32 Xuanwen Xu 36	Arith citrate aned as osteoblast-specific fact tein. It is present in a wide var velopment and function. Stud ncluding lung, ovary, breast an in the normal tissue and in the ce. The isoforms of periostin a rized by individual presence of 59). bmed ID Journal 964617 J Biomed Ma 197319 ACS Appl Ma 341935 Matrix Biol ter shipment. 9% glycerol, pH7.3	iety of normal adult tissues and fetal tiss ies show that periostin is overexpressed i nd colon cancers. Recent evidence reveals a stroma of the primary tumour, and it is r re between 83 and 93 kDa in mass and dit or absence of cassette exons 17-21 Application ther Res A IHC ater Interfaces IF	ues, a in a 5 that equire ffer in

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Immunohistochemical analysis of paraffinembedded human colon tissue slide using 19899-1-AP (Periostin antibody at dilution of 1:200 (under 10x lens). Immunohistochemical analysis of paraffinembedded human colon tissue slide using 19899-1-AP (Periostin antibody at dilution of 1:200 (under 40x lens).