

For Research Use Only

GLI3-Specific Polyclonal ANTIBODY



Catalog Number: 19949-1-AP

6 Publications

Basic Information

Catalog Number:

19949-1-AP

Size:

150UL, Concentration: 240 µg/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_000168

GeneID (NCBI):

2737

Full Name:

GLI family zinc finger 3

Calculated MW:

170 kDa

Observed MW:

190 kDa, 83-86 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:200-1:1000

IP 0.5-4.0 µg for IP and 1:200-1:1000 for WB

IHC 1:20-1:200

IF 1:10-1:100

Applications

Tested Applications:

FC, IF, IHC, IP, WB, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human placenta tissue, mouse lung tissue

IP: mouse lung tissue,

IHC: human testis tissue, human colon tissue

IF: HepG2 cells,

Background Information

GLI3 belongs to the GLI C2H2-type zinc-finger protein family. GLI3 plays a role in limb and brain development. GLI3 is implicated in the transduction of SHH signal. Defects in GLI3 are the cause of Greig cephalo-poly-syndactyly syndrome (GCPS). Defects in GLI3 are a cause of Pallister-Hall syndrome (PHS). Defects in GLI3 are a cause of type A1/B postaxial polydactyly (PAPA1/PAPB). Defects in GLI3 are a cause of type IV preaxial polydactyly. Defects in GLI3 are the cause of acrocallosal syndrome (ACS). The antibody is specific to GLI3. At the molecular level, Gli3 is translated into a 190-kDa transcriptional activator (Gli3-190) that undergoes proteolytic processing into a truncated 83-kDa repressor (Gli3-83) lacking C-terminal activation domains. (PMID: 16705181)

Notable Publications

Author	Pubmed ID	Journal	Application
Diana Trnski	26385428	Biochim Biophys Acta	WB
Yuqin Men	26549569	Sci Rep	WB
Maira H Nagai	32392211	PLoS Genet	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human placenta tissue were subjected to SDS PAGE followed by western blot with 19949-1-AP (GLI3-Specific antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of HepG2 cells using 19949-1-AP (GLI3-Specific antibody) at dilution of 1:25 and Rhodamine-Goat anti-Rabbit IgG.



IP Result of anti-GLI3-Specific (IP:19949-1-AP, 4ug; Detection:19949-1-AP 1:300) with mouse lung tissue lysate 4000ug.



Immunohistochemical analysis of paraffin-embedded human testis tissue slide using 19949-1-AP (GLI3-Specific Antibody) at dilution of 1:50.



1X10⁶ HepG2 cells were stained with 0.2ug GLI3-Specific antibody (19949-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.