For Research Use Only

NMDAR2A/GRIN2A Polyclonal antibody

Catalog Number:19953-1-AP 28 Publications




Basic Information	Catalog Number: 19953-1-AP	GenBank Accession Number: NM_000833	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 600 µg/ml by	2903	WB 1:500-1:1000	
	Nanodrop and 407 µg/ml by Bradford	UNIPROT ID:	IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	method using BSA as the standard;	Q12879	protein lysate	
	Source: Rabbit	Full Name:	IHC 1:200-1:800 IF/ICC 1:10-1:100	
	Isotype: IgG	glutamate receptor, ionotropic, N		
		methyl D-aspartate 2A Calculated MW:		
		165 kDa		
		Observed MW: 150-160 kDa		
Applications	Tested Applications:	Applications: Positive Controls:		
	WB, IP, IF, IHC, ELISA	WB: mouse brain tissue,		
	Cited Applications:	IP : mouse brain tissue,		
	WB, IP, IF		se brain tissue,	
	Species Specificity: human, mouse	IF/ICC : H		
	Cited Species:	in the comment		
	human, rat, mouse			
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen		
Background Information	GRIN2A is NMDA receptor subtype of	glutamate-gated ion channels pos gnesium. Activation of GRIN2A rec	gated ion channel (TC 1.A.10) family. sesses high calcium permeability and juires binding of agonist to both types c	
	GRIN2A is NMDA receptor subtype of voltage-dependent sensitivity to ma subunits. The antibody recognize the	glutamate-gated ion channels pos gnesium. Activation of GRIN2A rec		
	GRIN2A is NMDA receptor subtype of voltage-dependent sensitivity to ma subunits. The antibody recognize the Author Pu	glutamate-gated ion channels pos gnesium. Activation of GRIN2A rec C-term of GRIN2A.	sesses high calcium permeability and uires binding of agonist to both types o	
	GRIN2A is NMDA receptor subtype of voltage-dependent sensitivity to ma subunits. The antibody recognize the Author Pu Pengcheng Ma 36	glutamate-gated ion channels pos gnesium. Activation of GRIN2A rec C-term of GRIN2A. bmed ID Journal	sesses high calcium permeability and uires binding of agonist to both types o Application	
	GRIN2A is NMDA receptor subtype of voltage-dependent sensitivity to ma subunits. The antibody recognize theAuthorPuPengcheng Ma36Xin Peng34	glutamate-gated ion channels pos gnesium. Activation of GRIN2A rec C-term of GRIN2A. bmed ID Journal 179027 Sci Adv	sesses high calcium permeability and juires binding of agonist to both types o Application WB	
Background Information Notable Publications Storage	GRIN2A is NMDA receptor subtype of voltage-dependent sensitivity to ma subunits. The antibody recognize theAuthorPuPengcheng Ma36Xin Peng34	glutamate-gated ion channels pos gnesium. Activation of GRIN2A rec C-term of GRIN2A. bmed ID Journal 179027 Sci Adv 549339 J Mol Neurosci 876882 Sci Rep ter shipment.	sesses high calcium permeability and uires binding of agonist to both types o Application WB WB	


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



mouse brain tissue were subjected to SDS PAGE followed by western blot with 19953-1-AP (NMDAR2A/GRIN2A antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 19953-1-AP (NMDAR2A/GRIN2A antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 19953-1-AP (NMDAR2A/GRIN2A antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of Hela cells, using NMDAR2A/GRIN2A antibody 19953-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



IP result of anti-NMDAR2A/GRIN2A (IP:19953-1-AP, 4ug; Detection:19953-1-AP 1:600) with mouse brain tissue lysate 1680 ug.