For Research Use Only

OTUD5 Polyclonal antibody

Catalog Number:21002-1-AP 8 Publications



Basic Information

Catalog Number:

21002-1-AP

Rabbit

GenBank Accession Number:

BC028225

GeneID (NCBI):

Size: Genel D (N 150ul, Concentration: 600 ug/ml by 55593

Nanodrop;

UNIPROT ID: Q96G74 Full Name:

Isotype: OTU domain containing 5

IgG Calculated MW:
Immunogen Catalog Number: 571 aa, 61 kDa
AG15190 Observed MW:

70-75 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB: 1:500-1:3000

IF/ICC: 1:50-1:500

Applications

Tested Applications:

WB, IF/ICC, ELISA

Cited Applications: WB, IF, IP, CoIP

Species Specificity: human, mouse, rat

Cited Species: human, mouse **Positive Controls:**

WB: HEK-293T cells, HEK-293 cells, HeLa cells, Jurkat

cells, K-562 cells, NIH/3T3 cells, C6 cells

IF/ICC: A431 cells,

Background Information

The deubiquitinating enzyme OTUD5, also known as deubiquitinating enzyme A (DUBA), is a member of the deubiquitinating enzyme OTU family. By deubiquitinating other proteins, OTUD5 is an important regulator of several physiological processes, such as immune signaling and DNA damage responses. The dysregulation of these processes can increase the risk of associated diseases, including inflammation, cancer and genetic disorders.

Notable Publications

Author	Pubmed ID	Journal	Application
Qiangqiang Liu	35913115	Elife	WB
Junjing Zhang	40070026	Clin Transl Med	WB,IF
Xiaomeng Song	39721018	Adv Sci (Weinh)	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



HEK-293T cells were subjected to SDS PAGE followed by western blot with 21002-1-AP (OTUD5 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 21002-1-AP (OTUD5 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed A431 cells using OTUD5 antibody (21002-1-AP) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).