For Research Use Only

B3GNT6 Polyclonal antibody Catalog Number:21291-1-AP 5 Publications

proteintech® Antibodies | ELISA kits | Proteins www.ptglab.com


Basic Information	Catalog Number: 21291-1-AP Size: 150ul, Concentration: 600 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG15825	GenBank Accession Number: BC 103910	Purification Method: Antigen affinity purification		
		GenelD (NCBI):	Recommended Dilutions: WB 1:500-1:2000		
		192134			
		UNIPROTID: Q6ZMB0	IHC 1:50-1:500		
		Full Name: UDP-GlcNAc:betaGal beta-1,3-N- acetylglucosaminyltransferase 6 (core 3 synthase)			
				Calculated MW: 384 aa, 43 kDa	
				Observed MW:	
		43 kDa			
		Applications	Tested Applications:	Positive Controls: WB : mouse colon tissue, mouse small intestine tissue, mouse testis tissue	
			WB, IHC, ELISA		
Cited Applications: WB, IHC, IF					
Species Specificity: human, mouse					
Cited Species:					
human, mouse, rat					
Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0	vely, antigen				
Background Information	The B3GNT6 protein, also known as core 3 synthase, is a member of the O-GlcNAc transferase (OGT) family and add an N-acetylglucosamine to N-acetylgalactosamine-modifed serine or threonine. The B3GNT6 protein is responsible for the formation of the core 3 structure of O-glycans, which are important components of mucin-type glycoproteins This process is known as the protein O-GlcNAcylation modifcation. B3GNT6 expression has been reported to be largely downregulated in gastric and colorectal cancers. B3GNT6 has 3 isoforms with the molecular mass of 29 and 43 kDa. (PMID: 35387659)				
Notable Publications	Author Put	omed ID Journal	Application		
	Thomas G Biel 354	02630 Mol Ther Method	ds Clin Dev WB		
	J Ye 287	45318 Oncogene	WB,IHC		
	Bo Peng 374	55358 Int J Food Sci Nu	utr WB		
Storage	Storage: Store at -20°C. Stable for one year aft Storage Buffer: PBS with 0.02% sodium azide and 50	·			
	Aliquoting is unnecessary for -20 $^{\circ}$ C s	torage			
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	torage			

other manufacturer.

. W: ptglab.com

in USA), or 1(312) 455-8498 (outside USA)


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 21291-1-AP (B3GNT6 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 21291-1-AP (B3GNT6 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 21291-1-AP (B3GNT6 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 21291-1-AP (B3GNT6 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).