For Research Use Only

Integrin Alpha 3 Polyclonal antibody

Catalog Number:21992-1-AP

Featured Product

12 Publications



Basic Information

Catalog Number:

21992-1-AP

Nanodrop:

GenBank Accession Number:

BC150190

GeneID (NCBI):

Size: 150ul , Concentration: 500 ug/ml by

UNIPROT ID:

P26006 Rabbit Full Name:

Isotype: integrin, alpha 3 (antigen CD49C, alpha 3 subunit of VLA-3 receptor) IgG

Immunogen Catalog Number: Calculated MW:

AG17293 1066 aa. 119 kDa

> Observed MW: 130-140 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:5000-1:50000 IP: 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC: 1:50-1:500 IF/ICC: 1:500-1:2000

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity: human, mouse

Cited Species:

human, mouse

Positive Controls:

WB: A549 cells, human heart tissue, human testis

tissue, Calu-1 cells, PC-3 cells

IP: PC-3 cells.

IHC: mouse colon tissue, IF/ICC: A549 cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

ITGA3, also named as MSK18, CD49c, GAPB3 and FRP-2, belongs to the integrin alpha chain family. It is a receptor for fibronectin, laminin, collagen, epiligrin, thrombospondin and CSPG4. ITGA3 may mediate with LGALS3 the stimulation by CSPG4 of endothelial cells migration. ITGA3 is highly concentrated in epithelial cells where it strongly adheres to Laminin-5 and Laminin-5 induced rapid adhesion can be blocked by antibodies against the alpha-3 integrin subunit. It can be cleaved into heavy chain (840aa) and light chain (176aa). This antibody can recognize the heavy chain.

Notable Publications

Author	Pubmed ID	Journal	Application
Eunnyung Bae	35879332	Nat Commun	WB,IF
Qiqiao Du	32117712	Front Oncol	WB
Kexin Li	40118292	Biochim Biophys Acta Mol Basis Dis	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of Integrin Alpha 3 antibody (21992-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Integrin Alpha 3 transfected A549 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 21992-1-AP (Integrin Alpha 3 antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 21992-1-AP (Integrin Alpha 3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 21992-1-AP (Integrin Alpha 3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-Integrin Alpha 3 (IP:21992-1-AP, 4ug; Detection:21992-1-AP 1:500) with PC-3 cells lysate 1200ug.



Immunofluorescent analysis of (-20°C Methanol) fixed A549 cells using Integrin Alpha 3 antibody (21992-1-AP) at dilution of 1:1000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).