For Research Use Only

Basic Information

Glucocorticoid receptor Polyclonal antibody

Catalog Number:

Catalog Number:24050-1-AP

Featured Product

102 Publications

GenBank Accession Number:



Purification Method:

| Basic Information | 24050-1-AP | BC015610 | | Antigen affinity purification | |
|----------------------|---|--|--|--|---|
| | Size: 150ul, Concentration: 850 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG21146 | GeneID (NCBI): 2908 UNIPROT ID: P04150 Full Name: nuclear receptor subfa member 1 (glucocorti Calculated MW: 86 kDa Observed MW: 94-97 kDa | | Recommended Dilutions: WB 1:5000-1:50000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500 IF/ICC 1:200-1:800 | |
| Applications | Tested Applications: WB, IHC, IF/ICC, IP, ELISA | | | | |
| | Cited Applications: WB, IHC, IF, IP, CoIP, chIP Species Specificity: human, mouse, rat Cited Species: human, mouse, rat, chicken, zebrafish, rare minnow | | WB : HEK-293 cells, HeLa cells, HepG2 cells, DU 145 cells, mouse brain tissue, rat brain tissue, PC-12 cells IP : HepG2 cells, mouse heart tissue IHC : human prostate cancer tissue, human ovary | | |
| | | | | | |
| | | | Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 | | |
| | Background Information | receptor for glucocorticoids, which c response elements (GRE) and as a m | owns a dual mode of acti nodulator of other transc licated in newborn birth | on: as a transcri ription factors. I weight, thus pr | subfamily 3, group C, member 1) is a ption factor that binds to glucocorticoid t is involved in cell proliferation and oviding a biological mechanism by |
| Notable Publications | Author P | ubmed ID Jourr | nal | Application | |
| | Yixuan Wang 3 | 6165414 Br J P | harmacol | WB,IF,CoIP | |
| | Sheng Hu 3 | 6038029 Neuro | osci Lett | WB | |
| | Mehdi Boutagouga Boudjadja 3 | 6182699 Curr I | Biol | IF | |
| Storage | Storage: Stora at -20°C. Stable for one year a Storage Buffer: PBS with 0.02% sodium azide and 5 | i0% glycerol, pH7.3 | | | |


Aliquoting is unnecessary for -20 $^{\circ}$ C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 24050-1-AP (Glucocorticoid receptor antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 24050-1-AP (Glucocorticoid receptor antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



IP result of anti-Glucocorticoid receptor (IP:24050-1-AP, 4ug; Detection:24050-1-AP 1:4000) with HepG2 cells lysate 1720 ug.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Glucocorticoid receptor antibody (24050-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-Phalloidin (red).