

For Research Use Only

C10orf67 Polyclonal antibody

Catalog Number: 24459-1-AP



Basic Information

Catalog Number: 24459-1-AP	GenBank Accession Number: BC035732	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 750 µg/ml by Nanodrop and 347 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 256815	Recommended Dilutions: WB 1:200-1:1000
Source: Rabbit	UNIPROT ID: Q8IYJ2	
Isotype: IgG	Full Name: chromosome 10 open reading frame 67	
Immunogen Catalog Number: AG21465	Calculated MW: 184 aa, 21 kDa	
	Observed MW: 18-21 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : PC-3 cells, A549 cells, human testis tissue
Species Specificity: human	

Background Information

RNF17 also termed as ring finger protein 17 is a 1623 amino acid protein, which contains 1 ring type zinc finger and 4 tudor domains. RNF17 is specifically expressed in testis. The RING finger motif presents in many ubiquitin E3 ligases. RNF17 interacts with all four members of the Mad family (Mad1, Mxi1, Mad3 and Mad4), which are basic-helix-loop-helix-leucine zipper transcription factors of the Myc oncoprotein network. RNF17 is component of the tectonic-like complex, which is required for ciliogenesis and sonic hedgehog/SHH signaling. RNF17 is able to form dimers or polymers both in vitro and in vivo, indicating that it may play a role in the assembly of RNF17 granules. RNF17 encodes a novel key regulator of spermiogenesis.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



PC-3 cells were subjected to SDS PAGE followed by western blot with 24459-1-AP (C10orf67 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.