For Research Use Only

SLC25A22 Polyclonal antibody

Catalog Number:25402-1-AP 2 Publications


Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 25402-1-AP	GenBank Accession Nu BC023545	mber:	Purification Method: Antigen affinity purification	n
	Size:	GenelD (NCBI):		Recommended Dilutions:	
	150ul , Concentration: 1000 ug/ml by	79751		WB: 1:1000-1:6000	
	Nanodrop and 460 ug/ml by Bradford	UNIPROT ID: Q9H936		IHC: 1:50-1:500 IF-P: 1:50-1:500 IF/ICC: 1:50-1:500	
	method using BSA as the standard;				
	Source:	Full Name: solute carrier family 25 (mitochondrial carrier: glutamate), member 22		11/100.1.30-1.300	
	Rabbit				
	Isotype: IgG				
	Immunogen Catalog Number: AG19958	Calculated MW: 323 aa, 34 kDa Observed MW: 30-34 kDa			
Applications	Tested Applications:		Positive Controls:		
	WB, IHC, IF/ICC, IF-P, ELISA Cited Applications:		WB : mouse brain tissue, Caco-2 cells, HT-29 cells, I brain tissue, NCI-H1299 cells IHC : mouse brain tissue, human colon cancer tissue human liver cancer tissue		9 cells, ra
	WB Species Specificity:				er tissue,
	human, mouse, rat, monkey		IF-P : mouse brain tissue,		
	Cited Species: human, mouse		IF/ICC : COS-7 cells,		
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
	SLC25A22 encodes a mitochondrial glutamate/H+ symporter with highly expression in brain, especially in astrocytes, where it controls glutamate uptake. The mutations of SLC25A22 are associated with the early/infantile epileptic encephalopathies. It has been shown that SLC25A22 plays a key role in promoting proliferation and migration in cancer, such as human colorectal cancer (CRC) and Gallbladder cancer (GBC). And there is a result that the expression of SLC25A22 in GBC was significantly higher than that in adjacent tissues. 25402-1-AP antibody is specific to SLC25A22.(PMID: 30814911, 25033742, 24596948)				
Background Information	astrocytes, where it controls glutama epileptic encephalopathies. It has be migration in cancer, such as human c the expression of SLC25A22 in GBC w	te uptake. The mutation en shown that SLC25A22 olorectal cancer (CRC) a vas significantly higher t	s of SLC 25A 22 2 plays a key r nd Gallbladde :han that in ad	are associated with the early ole in promoting proliferation r cancer (GBC). And there is a	r/infantile n and result tha
Background Information	astrocytes, where it controls glutama epileptic encephalopathies. It has be migration in cancer, such as human c the expression of SLC25A22 in GBC w specific to SLC25A22.(PMID: 3081491	te uptake. The mutation en shown that SLC25A22 olorectal cancer (CRC) a vas significantly higher t	s of SLC25A22 2 plays a key r nd Gallbladde chan that in ad	are associated with the early ole in promoting proliferation r cancer (GBC). And there is a jacent tissues. 25402-1-AP ar	r/infantile n and result tha tibody is
	astrocytes, where it controls glutama epileptic encephalopathies. It has be migration in cancer, such as human c the expression of SLC25A22 in GBC v specific to SLC25A22.(PMID: 3081491	te uptake. The mutation en shown that SLC25A2: olorectal cancer (CRC) ar vas significantly higher t 1, 25033742, 24596948) bmed ID Journa	s of SLC25A22 2 plays a key r nd Gallbladde chan that in ad	are associated with the early ole in promoting proliferation r cancer (GBC). And there is a jacent tissues. 25402-1-AP ar Applice	r/infantile n and result tha tibody is
	astrocytes, where it controls glutama epileptic encephalopathies. It has be migration in cancer, such as human of the expression of SLC25A22 in GBC w specific to SLC25A22.(PMID: 3081491 Author Pu Maria F Noterman 33	te uptake. The mutation en shown that SLC25A22 olorectal cancer (CRC) an vas significantly higher t 1, 25033742, 24596948) bmed ID Journa 836587 Proc N	s of SLC25A22 2 plays a key r nd Gallbladde than that in ad	are associated with the early ole in promoting proliferation r cancer (GBC). And there is a jacent tissues. 25402-1-AP ar Applice	r/infantile n and result tha tibody is

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed COS-7 cells using 25402-1-AP (SLC25A22 antibody), at dilution of 1:200 and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L).




Various lysates were subjected to SDS PAGE followed by western blot with 25402-1-AP (SLC 25A 22 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 25402-1-AP (SLC 25A22 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 25402-1-AP (SLC 25A 22 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using SLC25A22 antibody (25402-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).