

For Research Use Only

EDA Polyclonal antibody

Catalog Number: 25892-1-AP

3 Publications



Basic Information

Catalog Number:

25892-1-AP

Size:

150ul, Concentration: 300 ug/ml by Nanodrop and 300 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG23154

GenBank Accession Number:

BC126143

GeneID (NCBI):

1896

UNIPROT ID:

Q92838

Full Name:

ectodysplasin A

Observed MW:

40 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:200-1:1000

IHC: 1:50-1:500

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC

Species Specificity:

human, mouse

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse liver tissue,

IHC : human liver tissue,

Notable Publications

Author	Pubmed ID	Journal	Application
Ye Ji Lee	39858559	Genes (Basel)	WB
Ya Zhang	39091933	Heliyon	IHC
Yuan Zhuang	38952411	Clin Cosmet Investig Dermatol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 25892-1-AP (EDA antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse liver tissue were subjected to SDS PAGE followed by western blot with 25892-1-AP (EDA Antibody) at dilution of 1:100 incubated at room temperature for 1.5 hours.