

For Research Use Only

TNF-alpha Polyclonal antibody

Catalog Number: 26405-1-AP

42 Publications



Basic Information

Catalog Number: 26405-1-AP	GenBank Accession Number: BC028148	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 500 ug/ml by Nanodrop;	GeneID (NCBI): 7124	Recommended Dilutions: WB: 1:500-1:2000 IHC: 1:50-1:500 IF/ICC: 1:50-1:500
Source: Rabbit	ENSEMBL Gene ID: ENSG00000232810	
Isotype: IgG	UNIPROT ID: P01375	
Immunogen Catalog Number: AG24020	Full Name: tumor necrosis factor (TNF superfamily, member 2)	
	Calculated MW: 233 aa, 26 kDa	
	Observed MW: 26 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA	Positive Controls: WB : THP-1 cells,
Cited Applications: WB, IHC, IF	IHC : human ovary tumor tissue, human tonsillitis tissue
Species Specificity: human	IF/ICC : PMA, LPS and Brefeldin A treated THP-1 cells,
Cited Species: human, mouse, rat, rabbit	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

TNF, as also known as TNF-alpha, or cachectin, is a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. It is expressed as a 26 kDa membrane bound protein and is then cleaved by TNF-alpha converting enzyme (TACE) to release the soluble 17 kDa monomer, which forms homotrimers in circulation. It is produced chiefly by activated macrophages, although it can be produced by many other cell types such as CD4+ lymphocytes, NK cells, neutrophils, mast cells, eosinophils, and neurons. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFR2. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, insulin resistance, and cancer.

Notable Publications

Author	Pubmed ID	Journal	Application
Yang Xu	35623314	Redox Biol	IHC
Yanling Wang	35914567	Acta Trop	IHC
Lihua Luo	35822247	Cell Prolif	IHC

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




PMA, LPS, and Brefeldin A treated THP-1 cells were subjected to SDS PAGE followed by western blot with 26405-1-AP (TNF Alpha antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with GAPDH Monoclonal antibody 60004-1-Ig as loading control.




Immunohistochemical analysis of paraffin-embedded human ovary tumor tissue slide using 26405-1-AP (TNF Alpha antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human ovary tumor tissue slide using 26405-1-AP (TNF Alpha antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed PMA, LPS and Brefeldin A treated THP-1 cells using TNF-alpha antibody (26405-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 26405-1-AP (TNF Alpha antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).