

For Research Use Only

# NDRG1 Polyclonal antibody

Catalog Number: 26902-1-AP

Featured Product

24 Publications



## Basic Information

### Catalog Number:

26902-1-AP

### Size:

150ul, Concentration: 950 ug/ml by Nanodrop and 467 ug/ml by Bradford method using BSA as the standard;

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG25424

### GenBank Accession Number:

BC006260

### GeneID (NCBI):

10397

### UNIPROT ID:

Q92597

### Full Name:

N-myc downstream regulated 1

### Calculated MW:

43 kDa

### Observed MW:

43 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB: 1:1000-1:4000

IHC: 1:400-1:1600

IF/ICC: 1:50-1:500

FC (Intra): 0.40 ug per 10<sup>6</sup> cells in a 100 µl suspension

FC: 0.40 ug per 10<sup>6</sup> cells in a 100 µl suspension

## Applications

### Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

### Cited Applications:

WB, IHC, IF, IP, CoIP

### Species Specificity:

human

### Cited Species:

human, mouse

### Positive Controls:

WB: HEK-293 cells, HeLa cells

IHC: human colon cancer tissue,

IF/ICC: HeLa cells,

FC (Intra): HeLa cells,

FC: HeLa cells,

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

NDRG1 is a member of the N-myc down-regulated gene family which belongs to the alpha/beta hydrolase superfamily. NDRG1 is a highly conserved cytoplasmic protein involved in cell growth arrest and proliferation, cell differentiation, DNA damage response, heavy metal response, the hypoxia response, tumorigenesis, and metastasis. NDRG1 is a potent metastasis suppressor in multiple tumor-types. NDRG1 appears to serve as a putative tumor suppressor and is down-regulated in colorectal, gastric, cervical, ovarian and prostate cancers.

## Notable Publications

Author	Pubmed ID	Journal	Application
Callum Tromans-Coia	34508355	Nucleic Acids Res	WB
Yuyang Liu	30922920	Exp Cell Res	WB, IHC
Elodie Villa	33756106	Mol Cell	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




HEK-293 cells were subjected to SDS PAGE followed by western blot with 26902-1-AP (NDRG1 Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




WB result of NDRG1 antibody (26902-1-AP; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NDRG1 transfected HEK-293 cells.




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 26902-1-AP (NDRG1 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 26902-1-AP (NDRG1 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using NDRG1 antibody (26902-1-AP) at dilution of 1:200 and CoraLite@488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



$1 \times 10^6$  HeLa cells were intracellularly stained with 0.4 ug Anti-Human NDRG1 (26902-1-AP)(red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).