

For Research Use Only

VWF Polyclonal antibody

Catalog Number: 27186-1-AP



Basic Information

Catalog Number:

27186-1-AP

Size:

150ul , Concentration: 600 µg/ml by Nanodrop and 333 µg/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG25578

GenBank Accession Number:

GeneID (NCBI):

7450

Full Name:

von Willebrand factor

Purification Method:

Antigen affinity purification

Recommended Dilutions:

IHC 1:50-1:500

IF 1:50-1:500

Applications

Tested Applications:

IF, IHC, ELISA

Species Specificity:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IHC : human tonsillitis tissue, rat brain tissue, mouse brain tissue

IF : human tonsillitis tissue,

Background Information

Von Willebrand factor (VWF) is a large multimeric glycoprotein found in blood plasma involved in hemostasis following vascular injury. Due to the multimeric nature of VWF, it can range in size from 500 to 20,000 kDa due to the differences in the number of subunits comprising the protein. Each subunit is approximately 250 kDa (PMID: 9759493). The biosynthesis of VWF in vivo is limited to endothelial cells (PMID: 4209883) and megakaryocytes (PMID: 2413071). VWF synthesized in endothelial cells is either released directly into the plasma via 27186a secretory pathway, or tubulized and stored in organelles unique to this cell type called Weibel-Palade bodies (PMID: 16459301). Whereas VWF synthesized in megakaryocytes is stored in the alpha granules of platelets (PMID: 2046403). The primary function of VWF is as an adhesive plasma glycoprotein, particularly factor VIII; an essential blood-clotting protein (PMID: 6982084). VWF is also important in platelet adhesion to wound sites by binding specifically to type I and type III collagen (PMID: 11098050), with larger VWF multimers being most effective (PMID: 24448155).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 27186-1-AP (VWF antibody) at dilution of 1:200 (under 10x lens). Proteolytic pre-treatment mediated antigen retrieval with .



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 27186-1-AP (VWF antibody) at dilution of 1:200 (under 40x lens). Proteolytic pre-treatment mediated antigen retrieval with .



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using 27186-1-AP (vWF antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).