For Research Use Only

TTC21B Polyclonal antibody

Catalog Number:30112-1-AP



Purification Method:

WB 1:2000-1:10000 IHC 1:400-1:1600

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number:

30112-1-AP NM_024753 GeneID (NCBI): Size: 150ul , Concentration: 650 ug/ml by 79809

Nanodrop: **UNIPROT ID:** Q7Z4L5 Rabbit Full Name:

Isotype: tetratricopeptide repeat domain 21B

IgG Calculated MW:

Immunogen Catalog Number: 151KD

AG32456 Observed MW:

55 kDa

Applications

Tested Applications:

WB, IHC, ELISA WB: A549 cells, HEK-293 cells, K-562 cells

Positive Controls:

IHC: mouse testis tissue,

Species Specificity: Human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

TTC21B, also known as THM1, IFT139, belongs to the TTC21 family, containing several tetratricopeptide repeat (TPR) domains. TTC21B is a component of the IFT complex A (IFT-A), a complex required for retrograde ciliary transport and entry into cilia of G protein-coupled receptors (GPCRs). TTC21B is essential for retrograde trafficking of IFT-1, IFT-B, and GPCRs (PubMed:27932497). TTC21B is localized to the cilium axoneme and may play a role in retrograde intraflagellar transport in cilia (PMID: 18327258). Mutations in TTC21B are associated with nephronophthisis 12 (NPHP12) and short-rib thoracic dysplasia 4 with or without polydactyly (SRTD4)(PMID: 21258341).

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


*** 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 30112-1-AP (TTC21B antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 30112-1-AP (TTC21B antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.