## For Research Use Only

## OGG1 Polyclonal antibody

Catalog Number: 30302-1-AP



**Purification Method:** 

WB 1:500-1:1000

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number: GenBank Accession Number:

30302-1-AP BC000657 GeneID (NCBI): Size:

150ul, Concentration: 500 ug/ml by 4968

Nanodrop; **UNIPROT ID:** 015527 Rabbit Full Name:

Isotype: 8-oxoguanine DNA glycosylase

IgG Calculated MW:

Immunogen Catalog Number: 22 kDa, 36-40 kDa, 45-57 kDa

AG32962 Observed MW:

36-40 kDa

**Applications** 

**Tested Applications:** 

WB, ELISA

Species Specificity:

human, rat

Positive Controls:

WB: A549 cells, rat skeletal muscle tissue

## **Background Information**

The DNA damages induced by ROS contain base modification, base loss, and DNA single strand breaks, which are usually repaired by the base excision repair (BER) pathway in both prokaryotes and eukaryotes. OGG1 (The human 8-oxoguanine glycosylase 1) is the primary enzyme in BER pathway, responsible for the excision of 7, 8-dihydro-8oxoguanine (8-oxoG), a mutagenic base byproduct that occurs as a result of exposure to reactive oxygen species. There's 8 isoforms of OGG1, with calculated MW 22 kDa, 36-40 kDa and 45-57 kDa. The difference among these isoforms is the C-terminal (317-345aa).

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30302-1-AP (OGG1 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.