For Research Use Only

SCAMP5 Polyclonal antibody

Catalog Number: 30328-1-AP



Purification Method:

WB 1:1000-1:8000

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number:

30328-1-AP BC024700
Size: GeneID (NCBI):
150ul , Concentration: 600 ug/ml by 192683

Nanodrop; UNIPROT ID:
Source: Q8TAC9
Rabbit Full Name:

Isotype: secretory carrier membrane protein 5

IgG Calculated MW:

Immunogen Catalog Number: 26 kDa

AG31837 Observed MW:

26 kDa

Applications

Tested Applications: Positive Controls:

WB; mouse brain tissue, mouse brain tissue (incubated

at 37℃)

Species Specificity: Human, mouse

Background Information

Secretory carrier-associated membrane protein 5 (SCAMP5) belongs to the to the SCAMP family. Secretory carrier membrane proteins (SCAMPs) constitute a group of membrane transport proteins in plants, insects and mammals. The mammalian genome contains five types of SCAMP genes, namely, SCAMP1-SCAMP5. (PMID:36217917, PMID:19234194). SCAMPs participate in the vesicle cycling fusion of vesicles and cell membranes and play roles in regulating exocytosis and endocytosis, activating synaptic function and transmitting nerve signals. Among these proteins, SCAMP5 is highly expressed in the brain and has direct or indirect effects on the function of the central nervous system (PMID:36217917). SCAMP5 regulates membrane transport, controls the exocytosis of synaptic vesicles and is related to secretion carrier and membrane function. In addition, SCAMP5 plays a major role in the normal maintenance of the physiological functions of nerve cells (PMID:36217917, PMID:33663553). For optimal WB detection of this membrane protein, we recommend to avoid boiling the sample after lysis.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30328-1-AP (SCAMP5 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.