

For Research Use Only

GDI1 Polyclonal ANTIBODY



Catalog Number: 55214-1-AP

Basic Information

Catalog Number: 55214-1-AP	GenBank Accession Number: NM_001493	Purification Method: Antigen affinity purification
Size: 150UL, Concentration: 387 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 2664	Recommended Dilutions: WB 1:500-1:2000
Source: Rabbit	Full Name: GDP dissociation inhibitor 1	
Isotype: IgG	Calculated MW: 51 kDa	
	Observed MW: 51-55 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : SH-SY5Y cells, human brain tissue, mouse brain tissue
Species Specificity: human, mouse, rat	

Background Information

GDP dissociation inhibitors (GDIs) are proteins that regulate the GDP-GTP exchange reaction of members of the Rab family. GDIs can bind and release GDP-bound Rab proteins from membranes. Two GDI proteins towards different Rab proteins have been identified. GDI1 interacts with almost all of the Rab proteins, while GDI2 interacts with Rab11 but not Rab3A. GDI1 is expressed primarily in neural and sensory tissues and also in secretory cells, displaying a diffuse, cytoplasmic distribution in cells. It runs as a 55kda protein in SDS-PAGE. (PMID: 7929030, PMID: 19570034). Defects in GDI1 are the cause of mental retardation X-linked type 41 (MRX41). Defects in GDI1 are the cause of mental retardation X-linked type 48 (MRX48). The antibody is specific to GDI1. And it has no cross reaction to GDI2.


Storage

Storage:
Store at -20°C.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



SH-SY5Y cells were subjected to SDS PAGE followed by western blot with 55214-1-AP (GDI1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.