

For Research Use Only

PGAP1 Polyclonal antibody

Catalog Number: 55392-1-AP

3 Publications



Basic Information

Catalog Number:

55392-1-AP

Size:

150ul, Concentration: 1000 ug/ml by Nanodrop and 353 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_024989

GeneID (NCBI):

80055

UNIPROT ID:

Q75T13

Full Name:

post-GPI attachment to proteins 1

Calculated MW:

105 kDa

Observed MW:

49-53 kDa, 85-90 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:20-1:200

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse

Cited Species:

human

Positive Controls:

WB : human testis tissue, HEK-293 cells, MCF-7 cells, mouse testis tissue

IHC : human testis tissue, human breast cancer tissue

IF/ICC : HeLa cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

PGAP1 belongs to the GPI inositol-deacylase family. It is involved in inositol deacylation of GPI-anchored proteins. GPI inositol deacylation may important for efficient transport of GPI-anchored proteins from the endoplasmic reticulum to the Golgi. PGAP1 is the GPI inositol-deacylase (PMID:14734546). This antibody is specific to PGAP1.

Notable Publications

Author	Pubmed ID	Journal	Application
Tatsuaki Kurosaki	36356584	Mol Cell	WB
Takahiro Nakane	30898876	J Biol Chem	WB
Yutong Wu	39291803	Cancer Med	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human testis tissue were subjected to SDS PAGE followed by western blot with 55392-1-AP (PGAP1 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human testis using 55392-1-AP (PGAP1 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human testis using 55392-1-AP (PGAP1 antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using PGAP1 antibody (55392-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).