For Research Use Only

IFITM1-Specific Monoclonal antibody Catalog Number:60074-1-Ig Featured Product 74 Publications




Basic Information	Catalog Number: 60074-1-lg	GenBank Accession Number: BC000897	Purification Method: Protein A purification	
	Size:	GenelD (NCBI):	CloneNo.:	
	150ul , Concentration: 2000 ug/ml by		5B5E2	
	Nanodrop;	UNIPROT ID:	Recommended Dilutions:	
	Source: Mouse Isotype: IgG2a Immunogen Catalog Number: AG2320	P13164 Full Name: interferon induced transmembrane protein 1 (9-27) Calculated MW: 14 kDa	WB 1:20000-1:100000	
			IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
			IHC 1:200-1:1000 IF/ICC 1:200-1:800	
			11/100 1.200 1.000	
		Observed MW: 14-17 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, IF/ICC, IP, ELISA		WB : TF-1 cells, K-562 cells, human testis tissue, HepG2 cells, HL-60 cells IP : HepG2 cells,	
	Cited Applications: WB, IHC, IF, IP, CoIP			
	Species Specificity:	IP : Hep		
	human	IHC : hu	IHC : human lymphoma tissue,	
	Cited Species: IF/ICC : K-562 cells, human, mouse			
	retrieval may be performed w	ith citrate		
	buffer pH 6.0			
Background Information	buffer pH 6.0 IFITM1(interferon induced transmemi belongs to the CD225 family. It has to mediates cellular innate immunity to virus (WNV), and dengue virus , by ind viral restriction factors that inhibit im	vo transmembrane domain and o at least three major human pa nibiting the early steps of replic fection mediated by the influen ents of the innate immune syste	thogens, influenza A H1N1 virus, West Nile ation. IFITM proteins are recently identified as za A virus (IAV) hemagglutinin (HA) protein. em to restrict HIV-1 infection. 60074-1-lg is a	
	buffer pH 6.0 IFITM1(interferon induced transmemi belongs to the CD225 family. It has to mediates cellular innate immunity to virus (WNV), and dengue virus, by ind viral restriction factors that inhibit in Also they serve as important compon mouse monoclonal antibody which sp	vo transmembrane domain and o at least three major human pa nibiting the early steps of replic fection mediated by the influen ents of the innate immune syste	serves as an IFN-induced antiviral protein that thogens, influenza A H1N1 virus, West Nile ation. IFITM proteins are recently identified as za A virus (IAV) hemagglutinin (HA) protein. em to restrict HIV-1 infection. 60074-1-lg is a	
	buffer pH 6.0 IFITM1(interferon induced transmembelongs to the CD225 family. It has to mediates cellular innate immunity to virus (WNV), and dengue virus, by inf viral restriction factors that inhibit in Also they serve as important compon mouse monoclonal antibody which sp Author Pul	vo transmembrane domain and b at least three major human pa- nibiting the early steps of replic fection mediated by the influen ents of the innate immune syst pecifically recognizing IFITM1 b	serves as an IFN-induced antiviral protein that thogens, influenza A H1N1 virus, West Nile ation. IFITM proteins are recently identified as iza A virus (IAV) hemagglutinin (HA) protein. em to restrict HIV-1 infection. 60074-1-lg is a but not IFITM2 or IFITM3.	
Background Information	buffer pH 6.0 IFITM1(interferon induced transmembelongs to the CD225 family. It has to mediates cellular innate immunity to virus (WNV), and dengue virus, by inl viral restriction factors that inhibit in Also they serve as important componemouse monoclonal antibody which spectrum and the server as a server and the server as a server as	vo transmembrane domain and o at least three major human pa- nibiting the early steps of replic fection mediated by the influen ents of the innate immune syste pecifically recognizing IFITM1 b perifically and Journal	serves as an IFN-induced antiviral protein that thogens, influenza A H1N1 virus, West Nile ation. IFITM proteins are recently identified as iza A virus (IAV) hemagglutinin (HA) protein. em to restrict HIV-1 infection. 60074-1-lg is a but not IFITM2 or IFITM3. Application	
	buffer pH 6.0 IFITM1(interferon induced transmembelongs to the CD225 family. It has to mediates cellular innate immunity to virus (WNV), and dengue virus, by ind viral restriction factors that inhibit in Also they serve as important componed mouse monoclonal antibody which spectrum and the serve as important componed by the serve as i	vo transmembrane domain and o at least three major human pa- nibiting the early steps of replic fection mediated by the influen ents of the innate immune syste pecifically recognizing IFITM1 b perifically recognizing IFITM1 b perifically Sci Rep	serves as an IFN-induced antiviral protein tha thogens, influenza A H1N1 virus, West Nile ation. IFITM proteins are recently identified as iza A virus (IAV) hemagglutinin (HA) protein. em to restrict HIV-1 infection. 60074-1-lg is a but not IFITM2 or IFITM3. Application WB	
	buffer pH 6.0 IFITM1(interferon induced transmembelongs to the CD225 family. It has to mediates cellular innate immunity to virus (WNV), and dengue virus, by ind viral restriction factors that inhibit in Also they serve as important componed mouse monoclonal antibody which spectrum and the serve as important componed by the serve as i	vo transmembrane domain and b at least three major human pa- nibiting the early steps of replic fection mediated by the influen ents of the innate immune syste becifically recognizing IFITM1 b comed ID Journal 266929 Sci Rep 256397 Viruses 537236 J Mol Biol er shipment. % glycerol, pH7.3	serves as an IFN-induced antiviral protein tha thogens, influenza A H1N1 virus, West Nile cation. IFITM proteins are recently identified as iza A virus (IAV) hemagglutinin (IA) protein. em to restrict HIV-1 infection. 60074-1-lg is a but not IFITM2 or IFITM3. Application WB WB	


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 60074-1-1g (IFITM1-Specific antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.




TF-1 cells were subjected to SDS PAGE followed by western blot with 60074-1-1g (IFITM1-Specific antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.




IP result of anti-IFITM1-Specific (IP:60074-1-Ig, 4ug; Detection:60074-1-Ig 1:500) with HepG2 cells lysate 3600ug.



Immunohistochemical analysis of paraffinembedded human lymphoma tissue slide using 60074-1-1g (IFITM1-Specific Antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lymphoma tissue slide using 60074-1-1g (IFITM1-Specific Antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed K-562 cells using IFITM1-Specific antibody (60074-1-Ig, Clone: 5B5E2) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).