For Research Use Only

B23/NPM1 Monoclonal antibody

Catalog Number:60096-1-lg Featured Product

44 Publications


oroteintech Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 60096-1-lg	GenBank Accession Number: BC002398		Purification Method: Protein G purification	
	Size:	GenelD (NCBI):		•	
	150ul , Concentration: 1000 ug/ml by			CloneNo.: 4F12A3	
	Nanodrop;			Recommended Dilutions:	
	Source:	P06748		WB: 1:5000-1:50000	
	Mouse	Full Name:		IHC: 1:200-1:1000	
	lsotype: lgG1	nucleophosmin (nucleolar IF/ICC: 1:200- phosphoprotein B23, numatrin) Calculated MW: 33 kDa Observed MW: 35-38 kDa		IF/ICC: 1:200-1:800	
	Immunogen Catalog Number: AG7415				
Applications	Tested Applications: WB, IHC, IF/ICC, ELISA		Positive Controls:		
		d Applications: LNCaP cel		B : HSC-T6 cells, C6 cells, HeLa cells, HepG2 cells, ICaP cells, NIH/3T3 cells, RAW 264.7 cells, ROS172 Ills, MCF-7 cells, HEK-293 cells, Jurkat cells, K-562 Ills IC : human breast cancer tissue, human colon cance	
	WB, IHC, IF, IP, ChIP				
	Species Specificity: cells				
			IHC : human		
	Cited Species:	tissue			
	human management		IF/ICC : Hep	G2 cells,	
	Note-IHC: suggested antigen re				
	TE buffer pH 9.0; (*) Alternation retrieval may be performed w buffer pH 6.0	ely, antigen			
Background Information	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0 Nucleophosmin (NPM1,B23) is a putate nuclear localization signals (NLSs). Th process involving several cytoplasmi pore complex and translocate across in nuclear proteins that are central to the	rely, antigen ith citrate ive ribosome assembly ne transport of proteins c factors. Proteins must the nuclear envelope ir s process have been id	across the nuc t be recognized n an ATP-depe lentified. The 3	lear envelope is a selective, multistep d as import substrates, dock at the nucle	
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0 Nucleophosmin (NPM1,B23) is a putate nuclear localization signals (NLSs). The process involving several cytoplasmis pore complex and translocate across is nuclear proteins that are central to thi involved in ribosomal assembly and kinase during mitosis.	rely, antigen ith citrate ive ribosome assembly ne transport of proteins c factors. Proteins must the nuclear envelope ir s process have been id	across the nuc t be recognized n an ATP-depe lentified. The 3 abundant prot	lear envelope is a selective, multistep d as import substrates, dock at the nucle ndent fashion. Several cytosolic and 8 kDa nuclear protein nucleophosmin is	
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0 Nucleophosmin (NPM1,B23) is a putat nuclear localization signals (NLSs). Th process involving several cytoplasmi process involving several cytoplasmi nuclear proteins that are central to thi involved in ribosomal assembly and kinase during mitosis.	rely, antigen ith citrate ive ribosome assembly the transport of proteins c factors. Proteins must the nuclear envelope ir s process have been id rRNA transport. It is an	across the nuc t be recognized n an ATP-depe lentified. The 3 abundant prot	lear envelope is a selective, multistep d as import substrates, dock at the nucle ndent fashion. Several cytosolic and 8 kDa nuclear protein nucleophosmin is ein that is highly phosphorylated by Cdo	
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0Nucleophosmin (NPM1,B23) is a putat nuclear localization signals (NLSs). Th process involving several cytoplasmi pore complex and translocate across to nuclear proteins that are central to thi involved in ribosomal assembly and kinase during mitosis.AuthorPub Shutong FanShutong Fan304	rely, antigen ith citrate ive ribosome assembly ne transport of proteins c factors. Proteins must the nuclear envelope in s process have been id rRNA transport. It is an med ID Journa 05806 Oncol	across the nuc t be recognized n an ATP-depe lentified. The 3 abundant prot	dear envelope is a selective, multistep d as import substrates, dock at the nucle- ndent fashion. Several cytosolic and 88 kDa nuclear protein nucleophosmin is ein that is highly phosphorylated by Cde Application	
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0Nucleophosmin (NPM1,B23) is a putate nuclear localization signals (NLSs). The process involving several cytoplasmin pore complex and translocate across in nuclear proteins that are central to the involved in ribosomal assembly and kinase during mitosis.AuthorPub Shutong FanShutong Fan304 245	rely, antigen ith citrate ive ribosome assembly ne transport of proteins c factors. Proteins must the nuclear envelope in s process have been id rRNA transport. It is an med ID Journa 05806 Oncol	across the nuc t be recognized n an ATP-depe lentified. The <u>3</u> abundant prot	lear envelope is a selective, multistep d as import substrates, dock at the nucle ndent fashion. Several cytosolic and 8 kDa nuclear protein nucleophosmin is ein that is highly phosphorylated by Cd Application IHC	
Background Information Notable Publications	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0Nucleophosmin (NPM1,B23) is a putate nuclear localization signals (NLSs). The process involving several cytoplasmin pore complex and translocate across in nuclear proteins that are central to the involved in ribosomal assembly and kinase during mitosis.AuthorPub Shutong FanShutong Fan304 245	rely, antigen ith citrate ive ribosome assembly ne transport of proteins c factors. Proteins must the nuclear envelope in s process have been id rRNA transport. It is an of 806 Oncol 51807 Mol Ne 34264 Brain er shipment.	across the nuc t be recognized n an ATP-depe lentified. The <u>3</u> abundant prot	lear envelope is a selective, multistep d as import substrates, dock at the nucle ndent fashion. Several cytosolic and 8 kDa nuclear protein nucleophosmin is ein that is highly phosphorylated by Cd Application IHC	


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 60096-1-1g (B23/NPM1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRPconjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.




WB result of B23/NPM1 antibody (60096-1-lg; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-B23/NPM1 transfected HepG2 cells.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60096-1-1g (B23 Antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60096-1-Ig (B23 Antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 60096-1-1g (B23/NPM1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using B23/NPM1 antibody (60096-1-lg, Clone: 4F 12A3) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).