For Research Use Only

HDAC4-specific Monoclonal antibody proteintech

Catalog Number:60105-1-lg 1 Publications



Basic Information

Catalog Number:

60105-1-lg

150ul , Concentration: 153 µg/ml by 9759 Bradford method using BSA as the

standard; Source:

Mouse Isotype: lgG1

Full Name: histone deacetylase 4 Calculated MW:

119 kDa

Applications

Tested Applications:

ELISA

Species Specificity:

human

Purification Method:

Caprylic Acid/Ammonium Sulfate

Precipitation

CloneNo.: 4C3B9

Background Information

HDAC4, also named as HDACA, belongs to the histone deacetylase family. HDAC4 is responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein $complexes. \ HDAC4 is Involved in muscle \ maturation \ via its interaction \ with \ the \ myocyte \ enhancer \ factors \ such \ as$ MEF2A, MEF2C and MEF2D. This antibody is a mouse monoclonal antibody raised against a peptide mapping within human HDAC4.

GenBank Accession Number:

BC039904

GeneID (NCBI):

UNIPROT ID:

P56524

Notable Publications

Author	Pubmed ID	Journal	Application
Natalia Sacilotto	27898394	Genes Dev	

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Selected Validation Data