For Research Use Only

FAM3C Monoclonal antibody

Catalog Number: 60282-1-Ig



Basic Information

Catalog Number: GenBank Accession Number:

60282-1-lg BC046932 GeneID (NCBI): Size:

150ul, Concentration: 1000 ug/ml by 10447 Nanodrop and 620 ug/ml by Bradford UNIPROT ID: method using BSA as the standard; Q92520

Source: Full Name: Mouse family with sequence similarity 3,

Isotype: member C lgG1 Calculated MW: 25 kDa Immunogen Catalog Number: AG5516 Observed MW:

25 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA

Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method: Protein G purification

CloneNo.: 2E10D5

Recommended Dilutions:

WB 1:500-1:2000 IHC 1:50-1:500 IF/ICC 1:200-1:800

Positive Controls:

WB: fetal human brain tissue,

IHC: human oesophagus cancer tissue,

IF/ICC: HepG2 cells,

Background Information

FAM3C, also known as ILEI, is a secreted protein belonging to the cytokine-like protein family. FAM3C is a signaling cytokine with a GG domain, hydrophobic leader sequence and an N-terminal signal peptide. A change in expression of this protein has been noted in pancreatic cancer-derived cells. FAM3C has been involved in the epithelial to mesenchymal transition, and retinal laminar formation processes in vertebrates. Also, FAM3C may play an important role in inner ear development and thus be involved in cell differentiation and proliferation during inner ear embryogenesis.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com


Selected Validation Data




fetal human brain tissue were subjected to SDS PAGE followed by western blot with 60282-1-1g (FAM3C Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human oesophagus cancer tissue slide using 60282-1-lg (FAM3C Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using FAM3C antibody (60282-1-Ig, Clone: 2E10D5) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1).



Immunohistochemical analysis of paraffinembedded human oesophagus cancer tissue slide using 60282-1-Ig (FAM3C Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).