For Research Use Only

Perilipin-2 Monoclonal antibody

Catalog Number: 60340-1-lg



Purification Method:

Protein G purification

CloneNo.:

Basic Information

Catalog Number: GenBank Accession Number:

60340-1-lg BC005127
Size: Genel D (NCBI):

 150ul , Concentration: 1000 ug/ml by
 123
 1G5D4

 Nanodrop;
 UNIPROT ID:
 Recommended Dilutions:

 Source:
 Q99541
 WB: 1:500-1:2000

 Mouse
 Full Name:
 IHC: 1:200-1:8000

Isotype: adipose differentiation-related

lgG1 protein

Immunogen Catalog Number: Calculated MW: AG7643 48 kDa

Observed MW: 47-52 kDa

Applications

Tested Applications: WB, IHC, ELISA

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: human adipose tissue,

IHC: human renal cell carcinoma tissue, mouse brown adipose tissue, rat brown adipose tissue

Background Information

ADRP (adipocyte differentiation related protein) also known as ADFP, adipophilin, or perilipin-2, is a member of PAT family which is responsible for the transportation of lipids and the formation of lipid droplets. ADRP is localized on the surface of lipid droplets in a variety of tissues and cell lines. ADRP is not detected in undifferentiated cells but increases rapidly to high levels when adipocyte precursors differentiate into adipocytes. Anti-ADRP antibody is a reliable and sensitive marker for lipid droplet. Enhanced expression of ADRP is linked to diseases with abnormal lipid storage, including hepatic steatosis, atherosclerosis and diabetes. Immunohistochemistry of ADRP may facilitate histomorphological diagnosis of these diseases.

Storage

Storage

Storage Ruffer:

Storage Buffe

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




human adipose tissue were subjected to SDS PAGE followed by western blot with 60340-1-lg (ADRP Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-Ig (ADRP Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-Ig (ADRP Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-1g (ADRP/Perilipin 2 antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-1g (ADRP/Perilipin 2 antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat brown adipose slide using 60340-1 Ig (ADRP/Perilipin 2 antibody) at dilution of 1:16000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).