For Research Use Only

Anti-Human CD117/c-Kit (104D2)

Catalog Number:65154-1-lg




Basic Information	Catalog Number: 65154-1-lg	GenBank Accession Number: BC071593	Purification Method:Purified by protein-A affinity chromatographyCloneNo.:104D2Recommended Dilutions:FC: 1.5 ug per 10^6 cells in 100 μl suspension	
	Size:	GeneID (NCBI): 3815 UNIPROT ID: P10721 Full Name: v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog		
	100ug , 1 mg/ml			
	Source: Mouse			
	Isotype:			
	IgG1, kappa			
		Calculated MW: 976 aa, 110 kDa		
Applications	Tested Applications: FC	Positive Co		
	Species Specificity: human, bovine	FC : TF-1 cells,		
Background Information	oncogene c-kit (PMID: 24481 CSF-1R, PDGFRβ, PDGFRα, ar progenitor cells, mast cells, melanocytes, germ cells, ast cells of Cajal) (PMID: 10582	CD117, also known as c-Kit and SCFR, is a transmembrane protein with tyrosine kinase activity encoded by the oncogene c-kit (PMID: 2448137). It is a member of the type III receptor tyrosine kinase family, which also includes CSF-1R, PDGFRB, PDGFRB, and FLT3 (PMID: 29518044). CD117 is expressed on hematopoietic stem cells and progenitor cells, mast cells, and is also found in a wide range of non-haemopoietic cell types (including melanocytes, germ cells, astrocytes, renal tubules, breast glandular epithelial cells, sweat glands, and interstitial cells of Cajal) (PMID: 10582338; 23073628). CD117 plays an important role in early haemopoiesis. It is also involved in pigmentation, fertility, gut movement, and some aspects of the nervous system (PMID: 23073628).		
Storage	Storage: Store at 2-8°C. Stable for one Storage Buffer: PBS with 0.09% sodium azid			

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^6 TF-1 cells were surface stained with 1.5 ug Anti-Human CD117 (65154-1-Ig, Clone:104D2) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (green), or stained with 1.5 ug Mouse IgG1 Isotype Control Antibody and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (black). Cells were not fixed.