For Research Use Only

14-3-3 Monoclonal antibody

Catalog Number:66061-1-lg 5 Publications



Purification Method:

CloneNo.:

2E10A3

Protein G purification

Recommended Dilutions:

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

WB: 1:2000-1:20000

protein lysate

IHC: 1:50-1:500

IF/ICC: 1:20-1:200

Basic Information

Applications

Catalog Number: GenBank Accession Number:

66061-1-lg BC056867 GeneID (NCBI):

150ul, Concentration: 1000 ug/ml by 10971 Nanodrop and 487 ug/ml by Bradford UNIPROT ID: method using BSA as the standard; P27348

Source: Full Name: Mouse tyrosine 3-

Isotype: monooxygenase/tryptophan 5lgG1 monooxygenase activation protein,

theta polypeptide Immunogen Catalog Number: AG6055 Calculated MW:

28 kDa Observed MW: 31 kDa, 28 kDa

Tested Applications:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

WB, IHC, IF/ICC, IP, ELISA WB: HEK-293 cells, HeLa cells, HepG2 cells, human Cited Applications: brain tissue, pig brain tissue, rat brain tissue, mouse WB, IHC, IF, IP, CoIP brain tissue

Positive Controls:

Species Specificity: IP: rat brain tissue, human, mouse, rat, pig

IHC: human lung cancer tissue, rat brain tissue Cited Species:

IF/ICC: HepG2 cells, human, mouse, rat, pig

buffer pH 6.0

14-3-3 proteins interact with a wide spectrum of proteins and possess diverse functions. Mammals express seven **Background Information** distinct 14-3-3 isoforms (gamma, epsilon, beta, zeta, sigma, theta, tau) that form multiple homo- and heterodimmers. 14-3-3 proteins display the highest expression levels in the brain, and have been implicated in several neurodegenerative diseases, including Alzheimer's disease and amyotrophic lateral sclerosis.

Notable Publications

Author	Pubmed ID	Journal	Application
Nerea Ugidos	31620119	Front Immunol	IF
Hui Zhou	33203800	Aging (Albany NY)	IHC
Yuetao Wen	35609730	Neurosci Res	WB,CoIP

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66061-1-1g (14-3-3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




IP result of anti-14-3-3 (IP:66061-1-1g, 4ug; Detection:66061-1-1g 1:20000) with rat brain tissue lysate 1080 ug.



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 66061-1-Ig (14-3-3 Antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 66061-1-Ig (14-3-3 Antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HepG2 cells using 66061-1-Ig (14-3-3 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).