

For Research Use Only

CD19 Monoclonal antibody

Catalog Number: 66298-1-Ig **3 Publications**



Basic Information

Catalog Number: 66298-1-Ig	GenBank Accession Number: BC006338	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 930	CloneNo.: 1C10A1
Source: Mouse	Full Name: CD19 molecule	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:500-1:2000
Isotype: IgG1	Calculated MW: 556 aa, 61 kDa	
Immunogen Catalog Number: AG19945	Observed MW: 95 kDa	

Applications

Tested Applications: FC, IHC, WB, ELISA	Positive Controls: WB : Raji cells, Ramos cells, Daudi cells IHC : human tonsillitis tissue,
Cited Applications: IHC, WB	
Species Specificity: human	
Cited Species: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

CD19 is a B cell-specific molecule that controls B cell activation by complexing with the BCR. CD19 is a member of the Ig superfamily and is the dominant component for the signaling complex on B cells that includes CD21, CD81, and CD22. Humans with CD19 deficiency have reduced proliferative responses to BCR stimulation in vitro and mount impaired Ab responses to vaccination.

Notable Publications

Author	Pubmed ID	Journal	Application
Xiaomin Lu	34700112	Int Immunopharmacol	IHC
Xinying Tang	32454143	Cancer Lett	WB
Runzhi Huang	33397394	Cancer Cell Int	IHC

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Raji cells were subjected to SDS PAGE followed by western blot with 66298-1-Ig (CD19 Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Untreated and PNGase F-treated lysates of Raji cells were subjected to SDS PAGE followed by western blot with 66298-1-Ig (CD19 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. PNGase F was obtained from Atagenix (cat.NO. ata808).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66298-1-Ig (CD19 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66298-1-Ig (CD19 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10⁶ Raji cells were stained with 0.20ug CD19 antibody (66298-1-Ig, red) and control antibody (blue). Fixed with 90% MeOH.