For Research Use Only

## NXN Monoclonal antibody

Catalog Number:66603-1-lg




Basic Information	Catalog Number: 66603-1-lg	GenBank Accession Number: BC009327	Purification Method: Protein G purification	
	Size: 150ul , Concentration: 2100 ug/ml by		CloneNo.: 2C6A4	
	Nanodrop and 1000 ug/ml by Bradfor method using BSA as the standard;	<sup>d</sup> UNIPROT ID: Q6DKJ4	Recommended Dilutions: WB 1:5000-1:50000	
	Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG9248	Full Name: nucleoredoxin	IHC 1:50-1:500 IF/ICC 1:400-1:1600	
		Calculated MW: 435 aa, 48 kDa		
		Observed MW: 48 kDa		
Applications	WB, IHC, IF/ICC, ELISA WB : He   Species Specificity: cells   human, mouse, rat IHC : hu   Note-IHC: suggested antigen retrieval with		e Controls:	
			WB : HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells IHC : human kidney tissue,	
			Background Information	Nucleoredoxin (NRX/NXN), a relatively uncharacterized member of the TRX family protein, has recently been reported to regulate the Wnt/β-catenin pathway, which itself regulates cell fate and early development, in a redox-dependent manner(PMID:17567240). Northern blot analysis detected several nucleoredoxin splice variants in adult mouse tissues, with highest expression in testis and skin and lowest expression in liver(PMID:9119370). It has 3 isoforms produced by alternative splicing. This antibody is specific to NXN.
Storage	Storage: Store at -20°C. Stable for one year aft Storage Buffer: PBS with 0.02% sodium azide and 50	% glycerol, pH7.3		
	Aliquoting is unnecessary for -20°C s			


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using NXN antibody (66603-1-Ig, Clone: 2C6A4) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 66603-1-Ig (NXN antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 66603-1-Ig (NXN antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 66603-1-1g (NXN antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.