## For Research Use Only

## MTAP Monoclonal antibody

Catalog Number:66706-1-Ig



**Basic Information** 

Catalog Number: GenBank Accession Number: 66706-1-lg BC018625

Protein G purification GeneID (NCBI): Size: CloneNo.:

150ul, Concentration: 1000 ug/ml by 4507 2B1G6

Nanodrop and 437 ug/ml by Bradford UNIPROT ID: Recommended Dilutions: method using BSA as the standard; Q13126 WB 1:2000-1:10000 Source: IHC 1:50-1:500 Full Name: methylthioadenosine phosphorylase IF/ICC 1:400-1:1600 Mouse

Isotype: Calculated MW: lgG1 283 aa, 31 kDa Immunogen Catalog Number: Observed MW: AG17954 31 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, ELISA

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: NIH/3T3 cells, HCT 116 cells, HSC-T6 cells, Raji

**Purification Method:** 

cells, HT-29 cells

IHC: mouse liver tissue, human kidney tissue, rat liver tissue, mouse kidney tissue, human colon cancer

tissue, human liver cancer tissue

IF/ICC: HeLa cells,

## **Background Information**

MTAP is a 5-deoxy-5-methylthioadenosine (MTA) phosphorylase, converting MTA to salvageable intermediates including adenine and 5-methylthioribose-1-phosphate. MTAP is abundant in normal cells, but it is frequently found to be deleted in a variety of cancers. Its deficiency is common in cancer cell lines as well as in primary leukemia, lung cancer, melanoma, bladder cancer, gliomas and breast cancer.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 66706-1-1g (MTAP antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 66706-1-Ig (MTAP antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using MTAP antibody (66706-1-1g, Clone: 2B1G6) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).