

For Research Use Only

ATG13

Monoclonal ANTIBODY



Catalog Number: 66708-1-Ig

Basic Information

Catalog Number: 66708-1-Ig	GenBank Accession Number: BC001331	Purification Method: Protein G purification
Size: 150UL, Concentration: 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 9776	CloneNo.: 1C3A7
Source: Mouse	Full Name: KIAA0652	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:50-1:500 IF 1:50-1:500
Isotype: IgG1	Calculated MW: 57 kDa	
Immunogen Catalog Number: AG12968	Observed MW: 57 kDa	

Applications

Tested Applications:

IF, IHC, WB, ELISA

Species Specificity:

Human, Mouse, Rat, Pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse brain tissue, BGC-823 cells, NIH/3T3 cells, pig brain tissue, rat brain tissue

IHC: mouse testis tissue, mouse brain tissue

IF: HEK-293 cells,

Background Information

ATG13 is one component protein of the ULK1 complex which is required for autophagosome formation and mitophagy. ATG13 has two nutrient regulatory phosphorylation sites and the phosphorylation status of ATG13 affect regulation of autophagy by modulating enzyme activity and cellular localization of ULK1. Besides, it has been reported the nonautophagic function of ATG13 on cardiac development for ATG13-deficient embryos show myocardial growth defects. (PMID:27387056, 26801615, 26644405)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




BGC-823 cells were subjected to SDS PAGE followed by western blot with 66708-1-Ig (KIAA0652 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 66708-1-Ig (KIAA0652 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



mouse brain tissue were subjected to SDS PAGE followed by western blot with 66708-1-Ig (KIAA0652 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 66708-1-Ig (KIAA0652 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using 66708-1-Ig (KIAA0652 antibody) at dilution of 1:100 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).